

6. Hill, D. P., Mooney, W. D., Fuis, G. S. and Walter, A., The Whispering Gallery in deep sedimentary basin (abstract). *Earthq. Notes*, 1982, **53**, 26.
7. Hwang, J. L. and Mooney, W. D., Velocity and Q -structure of the Central Valley, California, based on synthetic seismogram modeling of seismic refraction data. *Bull. Seismol. Soc. Am.*, 1986, **76**, 1053–1067.
8. King, D. and Falvey, M., A seismic survey in the Cooper Basin in Queensland. *Aust. Pet. Explor. Assoc. J.*, 1977, **17**(1), 78–84.
9. Lock, J. and Collins, C. D. N., Velocity depth modeling using refraction and reflection data recorded in the central Eromanya Basin, Queensland, Australia. *Tectonophysics*, 1983, **100**, 175–184.
10. McMechan, G. A. and Mooney, W. D., Asymptotic ray theory and synthetic seismograms for laterally varying structure, theory and application the Imperial Valley, California. *Bull. Seismol. Soc. Am.*, 1980, **70**, 2021–2035.
11. Prasad, A. S. S. R. S., Reddy, P. R. and Sarkar, D., Identification and usage of multiples in crustal seismics – An application in the Bengal basin, India. *Curr. Sci.*, 2002, **82**, 8–25.
12. Prasad, A. S. S. R. S., Reddy, P. R. and Sarkar, D., Use of multiples in construction of velocity structure, Deep Continental studies in India. *DST Newsletter*, 2003, **13**(1), 5–10.
13. Reddy, P. R., Prasad, A. S. S. R. S. and Sarkar, D., Velocity modeling of Bengal Basin refraction data – refinement using multiples. *J. Appl. Geophys.*, 1998, **39**, 109–120.
14. Biswas, S. K., Rift basins in western margin of India and their hydrocarbon prospects with special reference to Kutch basin. *Am. Assoc. Pet. Geol. Bull.*, 1982, **66**, 1497–1513.
15. Naqvi, S. M., *Geology and Evolution of the Indian Plate (From Hadean to Holocene – 4 Ga to 4 Ka)*, Capital Publishing Company, New Delhi, 2005, p. 450.
16. Prasad, G. V. R., Jaeger, J. J., Sahni, A., Gheerbrant, E. and Khajuria, C. K., Eutherian mammals from the Upper Cretaceous (Maastrichtian) inter-trappean beds of Naskal, Andhra Pradesh, India. *J. Vertebr. Paleontol.*, 1994, **14**, 260–277.
17. Vijaya Rao, V. and Tewari, H. C., Deep crustal seismic studies over the Indian Shield. In *Glimpses of Geoscience Research in India* (eds Singhvi, A. K., Bhattacharya, A. and Guha, S.), Angkor Publ., Noida, 2008, pp. 133–139.
18. Fuloria, R. C., Geology and hydrocarbon prospects of Mahanadi Basin, India. In (ed. Biswas, S. K.), Indian Petroleum Publishers, 1993. vol. 1, pp. 355–369.
19. Curray, J. R., Emmel, F. J., Moore, D. G. and Raitt, R. W., *Ocean Basin and Margins* (eds Nairn, A. E. M. and Stehli, F. G.), Plenum, New York, 1982, pp. 617–627.
20. Cerveny, V. and Pšencik, I., Program SEIS81: 2-Dimensional ray package, Research report, Institute of Geophysics, Charles University, Prague, 1981.
21. Tewari, H. C., The effect of thin high velocity layers on seismic refraction data: an example from Mahanadi Basin, India. *Pure Appl. Geophys.*, 1998, **15**, 63–79.
22. Tewari, H. C., Dixit, M. M. and Murty, P. R. K., Use of travel time skips in refraction analysis to delineate velocity inversion. *Geophys. Prospect.*, 1995, **43**, 793–804.
23. Reddy, P. R., Venkateswarlu, N., Prasad, A. S. S. R. S. and Koteswara Rao, P., Basement structure below the coastal belt of Krishna–Godavari Basin: correlation between seismic structure and well information. *Gondwana Res.*, 2002, **5**(2), 513–518.
24. Kaila, K. L., Tewari, H. C., Krishna, V. G., Dixit, M. M., Sarkar, D. and Reddy, M. S., Deep seismic sounding studies in the north Cambay and Sanchar basins, India. *Geophys. J. Int.*, 1990, **103**, 621–637.
25. Kaila, K. L., Reddy, P. R., Mall, D. M., Venkateswarlu, N., Krishna, V. G. and Prasad, A. S. S. R. S., Crustal structure of the West Bengal Basin, India, from deep seismic sounding investigations. *Geophys. J. Int.*, 1992, **111**, 45–66.
26. Rucker, M. L., Seismic refraction interpretation with velocity gradient and depth of investigation. In Proceedings of the Geophysics Conference, Los Angeles, California, 2002.

ACKNOWLEDGEMENTS. We thank the Director, NGRI, Hyderabad for permission to publish this work. We also thank Dr V. Vijaya Rao and Dr P. Prabhakara Prasad of NGRI for useful suggestions. Special thanks are due to anonymous reviewers.

Received 10 June 2010; revised accepted 19 October 2010

Elevated sea surface temperature during May 2010 induces mass bleaching of corals in the Andaman

P. Krishnan^{1,*}, S. Dam Roy¹, Grinson George¹, R. C. Srivastava¹, A. Anand², S. Murugesan¹, M. Kaliyamoorthy¹, N. Vikas³ and R. Soundararajan⁴

¹Marine Research Laboratory, Central Agricultural Research Institute, Port Blair 744 101, India

²Regional Remote Sensing Centre–Central, National Remote Sensing Centre (ISRO), Nagpur 440 010, India

³Dive India Ltd, Havelock, South Andaman 744 102, India

⁴No. 308/1, TNEB ‘A’ Flat, Belly Area, Anna Nagar, Chennai 600 040, India

Increasing sea surface temperature (SST) and its consequences on marine ecosystems are widely discussed. Andaman Sea witnessed a few bleaching events during 1998, 2002 and 2005. The present study was taken up to assess the extent of bleaching during 2010 in selected reef sites in the Andaman through line intercept transect survey. It was found that the fully bleached corals as a percentage of total coral cover were maximum at Havelock Island (69.49), followed by South Button Island (67.28), Nicolson Island (56.45), Red Skin Island (43.39), North Bay (41.65) and Chidiyatapu (36.54). Branching corals were the worst affected, whereas the massive corals were found to have relatively withstood the elevated SST. The status of reefs and the variability in bleaching with the progression of SST with respect to different coral species are discussed.

Keywords: Bleaching, climate change, coral reefs, sea surface temperature.

THE Andaman and Nicobar (A&N) Islands are bestowed with the richest coral diversity among all Indian reefs. A total of 177 species of hard corals falling under 57 genera have been reported from these islands¹. Recent surveys

*For correspondence. (e-mail: krishnanars@yahoo.com)

Table 1. Survey sites covered under the study

Study site	Latitude/longitude	Characteristics
Red Skin Island	11°32'23.27"N; 92°35'03.98"E	Within Mahatma Gandhi Marine National Park; open to tourists seasonally
North Bay	11°42'21.20"N; 92°45'15.20"E	Popular tourist site
Chidiyatapu	11°29'28.00"N; 92°42'40.00"E	Popular tourist site
Havelock–Aquarium	11°59'59.30"N; 92°55'38.00"E	Diving site
Havelock–Wall	12°03'26.60"N; 92°57'79.50"E	Popular diving site
South Button Island	12°13'42.20"N; 93°01'29.70"E	Protected island
Nicolson Island	12°04'52.00"N; 92°57'34.20"E	Protected island

Table 2. Biophysical status of reefs in the selected study sites (expressed as percentage)

Components of reef substrate	North Bay	Chidiyatapu	Red skin	Havelock–Aquarium	South Button	Havelock–Wall	Nicolson Island
Sand	10.32	42.36	54.09	14.17	0.00	0.00	0.00
Sand + dead corals	0.00	0.00	10.98	15.79	0.00	6.32	0.00
Dead corals	13.65	19.82	10.76	21.46	30.87	47.26	24.08
Dead corals + algae	0.00	0.00	1.48	0.00	0.00	0.00	0.00
Fully bleached corals	30.48	13.82	8.27	20.36	45.29	31.16	42.86
Partially bleached corals	41.59	22.91	8.72	19.11	12.12	13.68	24.39
Live corals (unbleached)	1.11	1.09	2.07	7.41	9.90	0.00	8.67
Soft corals (partially bleached)	2.86	0.00	3.16	1.01	1.83	1.58	0.00
Others (giant clam, anemone, etc.)	0.00	0.00	0.46	0.69	0.00	0.00	0.00
	100.00	100.00	100.00	100.00	100.00	100.00	100.00

have indicated that the numbers could be close to 80% of the global maximum². Bleaching is one of the major threats which has significantly affected the reefs across the globe during different time-periods^{3–6}. Corals have symbiotic association with zooxanthellae, the algae which are responsible for the colouration of the corals. Expulsion of the algae leads to the whitening of reef-building corals, widely referred to as bleaching. It is caused by physiological, algal, host-related stresses and various ecological and anthropogenic factors^{7–9}. Sea surface temperature (SST) is a critical factor for the wellbeing of symbiotic association of host animals like corals, giant clam and sea anemones with the microalgae.

Surveys conducted in the reefs of the Andaman reveal that the corals have been extensively bleached during April and May 2010 ranging from 37% to 70% in various sites. Similar bleaching events were reported in 1998 and 2002 in this region^{10–12}. However, the extent of the current bleaching surpasses the earlier observations.

The sites in the Andaman, which were surveyed under this pilot study during January–June 2010 are listed in Table 1. Three 100 m transects were laid for surveys on the bottom topography, live coral cover and extent of bleaching (full/partial)¹³. The atmospheric temperature and SST were measured *in situ* using a mercury bulb thermometer. Time-series data of air temperature of Andaman were obtained from the Automatic Weather Station with SD Data Logger (iMETOS, Australia), installed at Bloomsdale, South Andaman (11°38'50"N;

92°39'13"E). For the satellite-derived SST data, MODIS Global Level 3 Mapped Thermal IR daytime SST from Aqua and Terra sensors, available at the Physical Oceanography Distributed Active Archive Center (PODAAC) site of NASA was used¹⁴. Maps with 4.63 km spatial resolution covering the area surrounding the A&N Islands were analysed. Daily SST maps for the period 20 April to 12 May 2010 were analysed to see the change in SST during the period when bleaching was observed. Eight-day averaged SST maps from 4 April to 12 May of 2002 and 2005 were also studied. SST maps from the NASA JPL–PODAAC site available in HDF format were read using the software binary codes provided with the data and converted to tiff image format. Further processing was done using ERDAS–IMAGINE and ARC–GIS software for display and analysis of the maps. SST in the vicinity of the selected coral bleaching sites was noted for the period during which bleaching was observed.

The reef substratum in the selected sites was studied (Table 2) and the extent of live coral cover as a percentage of different components of reef substrates, excluding sand was found to be the highest in North Bay (81.60), followed by the Nicolson Island (75.92), South Button Island (67.31), Chidiyatapu (65.61) and Havelock Island (Wall and Aquarium; 45.80). The live corals as estimated here included both bleached and unbleached ones. Among the selected sites, the percentage cover of reef associates and soft corals was the highest in the transects laid in Red Skin island (7.89).




Figure 1. Temporal variation in maximum air temperature in Andaman.




Figure 2. Modis Level 3 SMI Aqua and Terra daytime sea surface temperature (SST) showing gradual increase from 20 April to 9 May 2010.

The bleaching of corals started in the first week of May 2010. The preceding month was characterized by hot and humid climate during which the maximum atmospheric and SST recorded were 34°C and 31°C respectively. During summer, SST rise of 2–3°C above the normal maximum can kill the corals¹⁵. Corals can also live in hotter parts where the summer temperature reaches 31°C (ref. 16). The average sea-water temperature during the first week of May in all the study sites was $33.8 \pm 0.6^\circ\text{C}$, which resulted in extensive bleaching in different parts of the islands. The atmospheric temperature showed a sudden increase in air temperature in the first, third and fourth weeks of April (Figure 1). The air temperature

remained high in the study sites up to the first week of May, with corresponding increase in SST. Subsequently, the temperature dropped to approximately 30°C, due to rains.

The MODIS-derived SST maps for selected time-periods for the areas surrounding A&N Islands (Figure 2) showed consistently high temperatures in the range 31–32°C during the last two weeks of April 2010 in the sites, except in Chidiyatapu, where it ranged from 32.1°C to 33.3°C. The temperature was seen to have increased during the first week of May to around 33°C until the second week of May and dropped with the onset of monsoon in the second–third weeks of May. As the A&N Islands are




Figure 3. Bleached corals as observed during May 2010. *a*, Branching coral (*Acropora* spp.); *b*, Plate coral (*Echinopora lamellosa*); *c*, Massive coral (*Porites solida*) and *d*, Partially bleached soft corals (*Simularis* sp.).

covered with clouds during most of the year, it is difficult to obtain continuous and cloud-free SST imagery. The average SST in the similar period during 2002 and 2005, when varying degrees (20–40%) of bleaching were observed in the Andaman^{10,11}, was $32.5 \pm 0.6^\circ\text{C}$ and $30.8 \pm 0.5^\circ\text{C}$ respectively. Eight-day SST data of 2002, when higher incidence of coral bleaching was recorded at these sites, showed that the temperature ranged from 31.6°C to 32.2°C in the last three weeks of April and moderated to 31.3°C – 31.9°C during the first week of May 2002. The SST at Chidiyatapu, however, showed higher values in the range 32.2 – 32.7°C during April 2002. The weekly SST for the last three weeks of April 2005 ranged between 30.8°C and 32.2°C , and it decreased to 29.8 – 31.1°C during the first week of May 2005. These observations clearly indicate the development of relatively warm water masses near and around the islands during the last three weeks of April almost every year. Vivekanandan *et al.*¹⁷ observed that bleaching occurred when the summer SST maxima exceeded 31°C and remained high for over three days. The current bleaching event, considered to be the worst since 1998, can be attributed to the prolonged higher SST ($> 31^\circ\text{C}$) in the region during April–May. It is predicted that the annual average SST would increase in all the regions in the Indian seas by 3 – 3.5°C during 2000–99, and in Andaman region, if the summer temperature exceeded

31.4°C for over a few weeks, then bleaching would occur¹⁷.

It was observed that the branching corals (*Acropora* spp.) were the worst affected due to bleaching. The predominant species were *Acropora formosa*, *Acropora nobilis*, *Acropora robusta*, *Acropora breuggemanni* and *Acropora grandis*. In South Buton, vast beds of *Acropora* spp. recorded almost 100% bleaching. In one of the dive spots in Havelock (Wall), the encrusting corals, *Diplostrea heliopora* were extensively bleached, whereas the plate corals, *Echinopora lamellosa* were the predominant fully bleached coral species in North Bay. Even massive corals (*Porites* sp.) were found to have fully bleached in some of the study sites, though the overall extent of bleaching was less compared to fragile branching corals. The variability in the impact of corals with respect to thermal stress has been already reported^{18–20}. It is important to note that while corals can survive acute short-term exposures over a few hours^{7,21}, there are no examples of coral species or genotypes that can survive chronic exposures to 3 – 6°C increase above the summer sea temperature for more than a month at a particular site^{22,23}. Hoegh-Guldberg *et al.*²⁴ observed that there could be variation in the onset of bleaching between species, but no species is adapted to sustain even a marginal increase in SST for a prolonged exposure. Mass bleaching of soft corals has been reported in Australia⁶ and Thailand²⁵. However, in




Figure 4. Status of actual coral cover in Andaman and extent of bleaching.




Figure 5. Reef associates affected by elevated SST during May 2010. *a*, Bleached sea anemone (*Heteractis magnifica*); *b*, Bleached giant clam (*Tridacna* sp.) and *c*, Withering brittle stars.

all the survey sites in the Andaman, soft corals belonging to *Sarcophyton* sp. and *Sinularia* sp. had been only partially bleached. The different types of corals bleached due to elevated SST are shown in Figure 3.

The coral cover estimated at the different study sites was analysed separately to determine the extent of bleaching. It was found that the fully bleached corals as a percentage of total estimated coral cover (unbleached, partially bleached and fully bleached) were maximum at Havelock Island (Wall; 69.49), followed by South Button Island (67.28), Nicolson Island (56.45), Havelock Island (43.45), Red Skin Island (43.39), North Bay (41.65) and Chidiyatapu (36.54). The percentage of normal corals (unaffected by the elevated SST) was maximum at the diving sites studied in Havelock Island (15.80), followed by South Button Island (14.71) and Red Skin Island (10.86, Figure 4). The bleaching was not confined to the

reef-building corals, but also observed among some of the reef communities like sea anemone and giant clam, which are also known to have symbiotic association with the zooxanthellae. The brittle stars in South Button Island were apparently in stress on the first day of observation of bleaching, but started withering their arms as the period of elevated SST prolonged and the extent of bleaching in corals progressed (Figure 5).

The fact that even 1–2°C increase in temperature results in widespread bleaching implied that most corals live close to their thermal limits²⁶. However, it was observed that there was variation in the bleaching pattern between the species and within the same species in different locations, which could be attributed to the higher thermal thresholds of certain species owing to their exposure to high temperature in their sub-habitats (e.g. warm tidal pools)²⁷. In the current study no variation in the



Figure 6. Corals in South Button Island showing filamentous algal growth over the fully bleached corals (August 2010).

coral-associated fishes could be found at the study sites following bleaching, as reported earlier¹⁶. The variation in the susceptibility of different coral species to elevated temperature would mean that climate change would result in different yet less diverse reef communities in the short term^{24,26}.

Bleaching and subsequent growth of new coral species is part of the natural selection process¹⁶. Indian coral reefs have experienced 29 widespread bleaching events since 1989 (www.reefbase.org), but intense bleaching occurred in 1998 and 2002 (refs 16, 28). The corals which were affected to the extent of 60–70% during the earlier events of bleaching recovered, but those fully bleached died. Subsequent surveys in August indicated that algae have deposited over the fully bleached corals (Figure 6). It is only hoped that the affected corals in the current bleaching event in the Andaman would recover to their original status, given the prevailing favourable environment, following the onset of monsoon. However, the frequency of bleaching events is set to increase¹⁷ owing to the global trends in the atmospheric and SST. Hence systematic studies on the status of the affected reefs in the Andaman would provide valuable clues on possible alteration in the thermal tolerance of corals in the region.

1. Venkataraman, K., Sathyanarayana, Ch., Alfred, J. R. B. and Wolstenhome, J., *Handbook on Hard Corals of India*, Zoological Survey of India, 2003, p. 266.
2. Turner, J. R. *et al.*, Report of the Phase I: Remote sensing and rapid site assessment survey, Coral reef systems of the Andaman Islands, GoI/UNDP GEF, 2001, p. 76.
3. Hoegh-Guldberg, O., Climate change, coral bleaching and the future of the world's coral reefs. *Mar. Freshwater Res.*, 1999, **50**, 839–866.
4. Celliers, L. and Schleyer, M. H., Coral bleaching on high-latitude marginal reefs at Sodwana Bay, South Africa. *Mar. Pollut. Bull.*, 2002, **44**, 1380–1387.
5. Whelan, K. R. T., Miller, J., Sanchez, O. and Patterson, M., Impact of the 2005 coral bleaching event on *Porites porites* and *Colpophyllia natans* at Tektite Reef, US Virgin Islands. *Coral Reefs*, 2007, **26**, 689–693.
6. Fabricius, K. E., Tissue loss and mortality in soft corals following mass-bleaching. *Coral Reefs*, 1999, **18**, 54.
7. Hoegh-Guldberg, O. and Smith, J. G., The effect of sudden changes in temperature, light and salinity on the population density and export

of zooxanthellae from the reef corals *Stylophora pistillata* and *Seriatopora hystrix*. *J. Exp. Mar. Biol. Ecol.*, 1989, **129**, 279–303.

8. Brown, B. E., Coral bleaching: causes and consequences. *Coral Reefs*, 1997, **16**, 129–138.
9. Fitt, W., Brown, B., Warner, M. and Dunne, R., Coral bleaching: interpretation of thermal tolerance limits and thermal thresholds in tropical corals. *Coral Reefs*, 2001, **20**, 51–65.
10. Soundararajan, R., Mass bleaching of coral reefs in the A and N Islands in 1998. *SANE Newsl.*, 2002.
11. DamRoy, S., Krishnan, P. and George, G., Assessment of coral reef health using satellite data. In Annual Report 2005–2006 (eds Srivastava, R. C. *et al.*), Central Agricultural Research Institute, Port Blair, 2006, pp. 73–74.
12. Pet-Soede, L., Wafar, M., Venkataraman, K., Rajan, P. T. and Wilhelmsson, D., The status of the coral reefs of India following the bleaching event of 1998. In *Coral Reef Degradation in the Indian Ocean*, Status Report 2000 (eds Souter, D., Obura, D. and Linden, O.), CORDIO, SAREC Marine Science Program, Stockholm University, Sweden, pp. 69–74.
13. English, S., Wilkinson, C. and Baker, V. (eds), *Survey Manual for Tropical Resources*, Australian Institute of Marine Sciences, Townsville, Australia, 1997, 2nd edn, p. 390.
14. PODAAC, 2007; http://podaac.jpl.nasa.gov:2031/DATASET_DOCS/modis_sst.html
15. Hyne, J., Bleaching, the great unknown. *Reef Management News. Reef Res.*, 1998, **8**, 8–11.
16. Kumaraguru, A. K., Jayakumar, K. and Ramakritinan, C. M., Coral bleaching 2002 in the Palk Bay, southeast coast of India. *Curr. Sci.*, 2003, **85**(12), 1787–1793.
17. Vivekanandan, E., Ali, M. H., Jasper, B. and Rajaopalan, M., Vulnerability of corals to warming of the Indian seas: a projection for the 21st century. *Curr. Sci.*, 2009, **97**(11), 1654–1657.
18. Hoegh-Guldberg, O. and Salvat, B., Periodic mass-bleaching and elevated sea temperatures – bleaching of outer reef slope communities in Moorea, French-Polynesia. *Mar. Ecol. Prog. Ser.*, 1995, **121**, 181–190.
19. Marshall, P. A. and Baird, A. H., Bleaching of corals on the Great Barrier Reef: differential susceptibilities among taxa. *Coral Reefs*, 2000, **19**, 155–163.
20. Loya, Y., Sakai, K., Yamasato, K., Nakano, Y., Sambali, H. and Van Woesik, R., Coral bleaching: the winners and the losers. *Ecol. Lett.*, 2001, **4**, 122–131.
21. Takahashi, S., Nakamura, T., Sakamizu, M., Woesik, R. and Yamasaki, H., Repair machinery of symbiotic photosynthesis as the primary target of heat stress for reef-building corals. *Jpn. Soc. Plant Physiol.*, 2004, **25**, 1–255.
22. Ulstrup, K. E., Berkelmans, R., Ralph, P. J. and Van Oppen, M. J. H., Variation in bleaching sensitivity of two coral species across a latitudinal gradient on the Great Barrier Reef: the role of zooxanthellae. *Mar. Ecol. Prog. Ser.*, 2006, **314**, 135–148.

23. Middlebrook, R., Hoegh-Guldberg, O. and Leggat, W., The effect of thermal history on the susceptibility of reef-building corals to thermal stress. *J. Exp. Biol.*, 2008, **211**, 1050–1056.
24. Hoegh-Guldberg, O. *et al.*, Coral reefs under rapid climate change and ocean acidification. *Science*, 2007, **318**, 1737–1742.
25. Chavanich, S., Viyakarn, V., Loyjiw, T., Pattaratamrong, P. and Chankong, A., Mass bleaching of soft coral, *Sarcophyton* spp. in Thailand and the role of temperature and salinity stress. *ICES J. Mar. Sci.*, 2009, **66**(7), 1515–1519.
26. Maynard, J. A., Anthony, K. R. N., Marshall, P. A. and Masiri, I., Major bleaching events can lead to increased thermal tolerance in corals. *Mar. Biol.*, 2008, **155**, 173–182.
27. Hoegh-Guldberg, O., Climate change and coral reefs: Trojan horse or false prophecy? *Coral Reefs*, 2008, **28**, 569–575.
28. Arthur, R., Coral bleaching and mortality in three Indian reef regions during an El Niño southern oscillation event. *Curr. Sci.*, 2000, **79**, 1723–1729.

Received 30 June 2010; revised accepted 22 October 2010

Molecular taxonomy of marine mammals stranded along Kerala coast, India

Sanil George¹, K. Meenakshi¹ and A. Bijukumar^{2,*}

¹Chemical Biology Group, Rajiv Gandhi Centre for Biotechnology, Thiruvananthapuram 695 014, India

²Department of Aquatic Biology and Fisheries, University of Kerala, Thiruvananthapuram 695 581, India

Application of molecular tools for the identification of threatened marine mammals has gained importance in recent years. Though live and dead strandings of cetaceans are common along the Indian coasts, the specimens are often not properly identified due to the lack of local taxonomic expertise and poor quality of the specimens. Two marine mammals washed ashore in a putrefied condition at Edayar (08°25'N lat., 76°57'E long.), Thiruvananthapuram District, southwest coast of India, were identified by sequencing of 16S rRNA and COI genes. Sequence and phylogenetic similarity search done with all entries in the DNA sequence database, GenBank using BLAST identified the stranded mammals as Bryde's whale (*Balaenoptera edeni*) and finless porpoise (*Neophocaena phocaenoides*). The present report is the second record of *B. edeni* from the southwest coast of India.

Keywords: Cetaceans, molecular taxonomy, phylogeny, stranded mammals.

THE marine mammal diversity of Indian seas, represented by around 30 recorded species, forms almost one-fourth of the world's marine mammals, and almost 8% of all mammalian fauna recorded in India¹. The qualitative

deficiency of data on marine mammals in India notwithstanding, the shortcomings of marine mammal research in India include geographic disproportionateness in records, non-reporting of mortality due to fishing operations, lack of peer review, incorrect identification of species, incorrect geographic information, inaccuracy in measurements, repeated citation of incorrect records, misinterpretation of observations and lack of molecular data². Though live and dead strandings of marine mammals, especially cetaceans (whales, dolphins and porpoises), are common along the Indian coasts, the specimens are often not properly identified due to the lack of local taxonomic expertise and poor condition of the specimens. Since all the cetaceans are of importance from the conservation point of view, documenting their presence in the ecosystem and precise taxonomy would provide valuable information on various aspects of distribution and migratory nature of different species in the seas around India.

The application of molecular techniques has given stimulating impulses to marine mammal identification. In particular, partial or complete sequences of mitochondrial rRNA genes have been evaluated for appropriate evolutionary rates to resolve some aspects of the higher groups, such as genus, family and order. Because they contain information from old splitting events in their conserved regions, fast-evolving parts should be useful to resolve more recent events, e.g. intraspecific or intra-generic³. In addition, DNA sequences of mitochondrial cytochrome oxidase subunit I (COI) gene have been used to estimate phylogenetic relationships among closely related species^{4,5}. DNA sequencing technology has provided us the ability to determine the source of tissue samples believed to be derived from threatened or endangered species⁶. A phylogenetic approach to identifying marine mammal sequences from unknown sources has gained support in recent years⁷, especially for identifying whale meat products in the open market^{6,8}. Above all, sequencing of mitochondrial DNA has been used for describing new species of marine mammals^{9–11} and for describing their phylogeny¹². From India, initial efforts have been made by the Central Marine Fisheries Research Institute, Cochin towards molecular identification of marine mammals^{13–15}.

Two putrefied carcasses of marine mammals were washed ashore at Edayar (08°25'N lat., 76°57'E long.), Thiruvananthapuram District, southwest coast of India on 27 June 2009 (Figure 1 a and b). Based on field observations, though the presence of long ventral pleats on the lower profile of the head extending up to the navel confirmed the identity of one specimen as a whale belonging to the genus *Balaenoptera* (Bryde's whale, fin whale and blue whale in this group, recorded from India)¹⁶, confusion prevailed in the identification because of the dented upper jaw. Further, detailed examination of the specimen (total length = 3.9 m) also was difficult since the body was rotten, emanating foul odour, and the demand from

*For correspondence. (e-mail: abiju@rediffmail.com)