

- cell entry of dsRNA to induce RNAi silencing. *Nature Cell Biol.*, 2006, **8**, 793–802.
27. Castanotto, D. and Rossi, J. J., The promises and pitfalls of RNA-interference-based therapeutics. *Nature*, 2009, **457**, 426–433.
 28. Kumar, M., Gupta, G. P. and Rajam, M. V., Silencing of acetylcholinesterase gene of *Helicoverpa armigera* by siRNA affects larval growth and its life cycle. *J. Insect. Physiol.*, 2009, **55**, 273–278.
 29. Kurreck, J., RNA interference: from basic research to therapeutic applications. *Angew. Chem.*, 2009, **48**, 1378–1398.
 30. Turner, C. T., Davy, M. W., MacDiarmid, R. M., Plummer, K. M., Birch, N. P. and Newcomb, R. D., RNA interference in the light brown apple moth, *Epiphyas postvittana* (Walker) induced by double-stranded RNA feeding. *Insect. Mol. Biol.*, 2006, **15**, 383–391.
 31. Griebler, M., Westerlund, S. A., Hoffmann, K. H. and Meyering-Vos, M., RNA interference with the allatregulating neuropeptide genes from the fall armyworm *Spodoptera frugiperda* and its effects on the JH titer in the hemolymph. *J. Insect. Physiol.*, 2008, **54**, 997–1007.
 32. Rourke, I. J. and East, P. D., Evidence for gene conversion between tandemly duplicated cytoplasmic actin genes of *Helicoverpa armigera* (Lepidoptera: Noctuidae). *J. Mol. Evol.*, 1997, **44**, 169–177.
 33. Wee, C. W., Lee, S. F., Robin, C. and Heckel, D. G., Identification of candidate genes for fenvalerate resistance in *Helicoverpa armigera* using cDNA-AFLP. *Insect Mol. Biol.*, 2008, **17**, 351–360.
 34. Zhang, Y. L. *et al.*, Identification, characterization, and expression of a novel *P450* gene encoding CYP6AE25 from the Asian corn borer, *Ostrinia furnacalis*. *J. Insect. Sci.*, 2011, **11**, 37.
 35. Campbell, P. M., Cao, A. T., Hines, E. R., East, P. D. and Gordon, K. H., Proteomic analysis of the peritrophic matrix from the gut of the caterpillar, *Helicoverpa armigera*. *Insect Biochem. Mol. Biol.*, 2008, **38**, 950–958.
 36. Bown, D. P., Wilkinson, H. S. and Gatehouse, J. A., Differentially regulated inhibitor-sensitive and insensitive protease genes from the phytophagous insect pest, *Helicoverpa armigera*, are members of complex multigene families. *Insect Biochem. Mol. Biol.*, 1997, **27**, 625–638.
 37. Carmen, S. M. and Juan, A. G., Antiviral strategies in plants based on RNA silencing. *Biochem. Biophys. Acta*, 2011, **1809**, 722–731.
 38. Nunes, F. M. F. and Simões, Z. L. P., A non-invasive method for silencing gene transcription in honeybees maintained under natural conditions. *Insect Biochem. Mol. Biol.*, 2009, **39**, 157–160.
 39. Zhou, X. G., Wheeler, M. M., Oi, F. M. and Scharf, M. E., RNA interference in the termite *Reticulitermes flavipes* through ingestion of double-stranded RNA. *Insect. Biochem. Mol. Biol.*, 2008, **38**, 805–815.
 40. Naito, Y., Yamada, T., Matsumiya, T., Ui-Tei, K., Saigo, K. and Morishita, S., dsCheck: highly sensitive off-target search software for dsRNA-mediated RNA interference. *Nucleic Acids Res.*, 2005, **33**, 589–591.

ACKNOWLEDGEMENTS. We thank ICAR, New Delhi for funding the subproject on 'Potential of RNAi in insect pest management: A model in silencing genes specific to tomato fruit borer, *Helicoverpa armigera* Hubner (Noctuidae: Lepidoptera)' under NAIP. We also thank to Dr Bengali Baboo, National Director, NAIP-PIU, New Delhi and Dr Sudhir Kochhar, National Coordinator (Component-4 – Basic & Strategic Research), NAIP-PIU, New Delhi for encouragement and Dr Amrik Singh Sidhu, Director, IIHR, Bangalore for encouragement and the necessary facilities. We gratefully acknowledge Dr K. R. Kranthi and Dr Sandhya Kranthi, CICR, Nagpur for providing populations of *H. armigera*.

Received 26 August 2011; revised accepted 15 May 2012

Consequences of underestimating ancient deforestation in South India for global assessments of climatic change

Gérard Bourgeon¹, K. M. Nair², B. R. Ramesh³ and Danny Lo Seen^{4,*}

¹3 rue du Guesclin, F-34000 Montpellier, France

²National Bureau of Soil Survey and Land Use Planning, Hebbal, Bangalore 560 024, India

³French Institute of Pondicherry, PO Box 33, Puducherry 605 001, India

⁴CIRAD-TETIS Joint Research Unit (UMR), F-34093 Montpellier Cedex 5, France

Land-cover changes occurring before 1800 are often ignored in the estimation of CO₂ emissions, probably because they are poorly documented in most tropical countries. India appears to be an exception to this rule. It was possible to reconstitute the main stages of the land-cover history for a large region of South India, and therefore to retrace the dynamics of CO₂ emissions during nearly 1000 years. It was then possible to demonstrate that 25% of the total emissions occurred before 1800, and are mistakenly considered as more recent emissions.

Keywords: Ancient deforestation, carbon dioxide emissions, climate change, global assessments, land cover.

IN 2004, deforestation and biomass decomposition accounted for 17.3% of the total 49 Pg of CO₂-equivalent anthropogenic emissions of greenhouse gases (GHGs)¹, while fossil fuel consumption represented 56.6%. Global estimates of recent emissions are computed from country-level inventories following IPCC guidelines^{2,3}. For past emissions, ancient land-cover changes are either neglected⁴ or introduced through a historical frame^{5,6}. In this communication, we assess the consequences of not considering ancient land-cover changes when estimating GHG anthropogenic emissions for a large study area located in South India (Figure 1), focusing on soil organic carbon (SOC) which represents the largest terrestrial C pool⁷.

In 2001, the Intergovernmental Panel on Climate Change (IPCC)⁴ calculations for land-use (LU) change related emissions amounted to 121 PgC for the period 1850–1990, with LU changes obtained by the difference between potential⁸ and current⁹ vegetation maps. Potential vegetation was represented by the 1850 land cover, with the implicit hypothesis that no significant deforestation occurred before that date, and that GHGs emissions due to land-use/land-cover changes had started at about the same time as those due to fossil-fuel consumption.

*For correspondence. (e-mail: danny.lo_seen@cirad.fr)

When Houghton and Hackler⁵, and Van Minnen *et al.*⁶ calculated emissions for tropical Asia and for India respectively, historical frames of land-cover changes were introduced. Houghton and Hackler⁵ considered the period before 1750 as the 'pre-disturbance period' (i.e. hypothesizing that no major anthropogenic land-cover changes had occurred before 1750), whereas Van Minnen *et al.*⁶ proposed four world maps of the 'reconstructed agricultural area (cropland and pasture)' for AD 1700, 1800, 1900 and 2000. In both studies, ancient land-cover changes were not considered, while in the latter one, the 1700s map⁶ surprisingly showed South India free of cultivated areas.

For estimating past SOC changes in our study area, reference stock (i.e. SOC stock under potential vegetation) was based on a potential vegetation map derived from the forest maps of Pascal *et al.*^{10,11} and Ramesh *et al.*¹². In these maps, the potential area of each vegetation type is mainly given by the bioclimate¹³. The potential vegetation types are either directly indicated on the forest maps or, in places where forests have been cleared, deduced from bioclimate maps¹³. According to the 'climax theory', this potential vegetation corresponds to the 'plesioclimax: i.e. the stage that would be reached within nearly a century if man and his cattle were to disappear'¹⁴.

The multifactor method used in previous studies^{15,16} for estimating the 1977 and 1999 SOC stocks was again used to compute the potential SOC stock, with potential

vegetation replacing the 1977 or 1999 land cover. A figure of 0.554 PgC was obtained for the potential SOC stock of the study area. In the same area, recent findings¹⁶ have established that the SOC stock was stable, around 0.433 PgC during the period 1977–1999 (0.4288 and 0.4369 PgC in 1977 (ref. 16) and 1999 (refs 15 and 16) respectively), and this despite an unabated deforestation. It was also established that the soil carbon losses in deforested areas were compensated by C sequestration elsewhere, mainly in recent plantations and in the newly irrigated croplands¹⁶. These explanations, which concern the SOC stock, can be probably extended to the biomass carbon stock because both are linked in the functioning of terrestrial ecosystems.

When comparing the potential SOC stock (0.554 PgC) to the 1999 stock (0.437 PgC), the difference, 0.117 PgC, represents an estimate of the overall loss of C from the soil, i.e. only 21.1% of the potential SOC stock.

We agree with authors^{5,6} who consider that the history of land-cover changes is key for understanding carbon stock variations. To integrate historical data in our ecological approach, we referred to the four historical modes of resources use (viz. gathering, nomadic pastoralism, settled cultivation and industry) defined by Gadgil and Guha¹⁷ to describe the ecological history of India. Then, in order to go beyond a simple qualitative description, we used the broad correspondences established by Ellis and Ramankutty¹⁸ between the main forms of human–ecosystem interaction and population density. The gathering mode corresponds to a population density < 1 person km⁻², while settled cultivation, and particularly irrigated cultivation, seems to need >100 persons km⁻². Intermediate densities (between 1 and 100 persons km⁻²) correspond to the other modes of resources use (shifting cultivation and nomadic pastoralism). We may notice that, in the type of irrigated agriculture in semi-arid India known as 'tank irrigation', these figures are in agreement with those of Van Oppen¹⁹, who showed that tanks appear when population density reaches 50–60 persons km⁻² and disappear when it exceeds 220 persons km⁻².

In India, archaeological records suggest that the southern part of the peninsula was occupied later, compared to the Indus Valley, and was also less intensively studied by archaeologists and historians²⁰. The period around 2500 BP was described as the starting time of settled agriculture and pastoralism in the drier parts of India¹⁷, after a long period when gathering was the only mode of resources use. Pastoralism is attested by numerous dung mounds in semi-arid Karnataka, but Fuller²¹ considered that most of these dung mounds were linked to seasonal encampments of a pastoral segment of the society rather than to permanent settlements. At the same period, because of the limited efficiency of stone-tools, the Western Ghats, as many other regions covered by moist forests, might have remained under the food-gathering mode associated with limited shifting cultivation¹⁷.




Figure 1. Location map showing the study area and the different places cited in the text.




Figure 2. Map of the Belgaum–Goa region, north of the study area. *a*, Corrected map of Gentil (1770) with Sarkars (e.g. Bankapour) and Parganas (e.g. Onar), the two lower levels of the Mughal administrative divisions; dotted lines represent boundaries of Sarkars. *b*, Modern map showing present-day districts.

The oldest iron tools discovered in Karnataka were dated from 1050 to 950 BC; this age corresponds to the ‘later Vedic period’ (1000–600 BC) of Randhawa²². The use of iron tools, axes, sickles and ploughs of different types allowed efficient deforestation and, at the same time, extension of ploughed areas and the true transition to settled agriculture.

Further agricultural development of South India was made by rulers of local kingdoms. The first one, the Chera kingdom, corresponding to the present-day Kerala, flourished during the second century AD (ref. 22). Other kingdoms appeared later, mostly to the west of the study area on the deltaic formations of the west coast, and have developed irrigated rice cultivation. Their rulers have continued on the same basis, encouraging land clearance and promoting tank and canal irrigation: Cholas in the present-day Tamil Nadu and Hoysalas in northeast Karnataka (Mysore–Hassan districts) for nearly three centuries (AD 1000–1300). The Hoysalas established their first capital in Halebidu (25 km north of Hassan, Figure 1) during the 11th century; they encouraged agriculture, forest clearance to extend cultivated lands, and promoted tank and canal irrigation²³.

The Portuguese were the first Europeans to settle in India during the 16th century (Cochin, Goa) introducing groundnuts, tobacco, potatoes and red chillies.

Later, the 18th century situation attracts particular attention because it corresponds approximately to the period chosen by many authors^{4–6} as the starting point of anthropogenic disturbance.

The maps of Gentil (1770) were helpful for documenting the 1750–1770s situation. Reprints of these historical maps were recently made available by Gole²⁴. On these maps, place names have been given and are ordered according to three administrative sub-divisions in use in the Mughal Empire at that time, viz. Subas, Sarkars and Parganas. The Parganas were places of administrative importance where taxes were collected. For comparison with more recent maps, the maps of Gentil had to be geometrically corrected because they were established from route marches without accurate topographic survey²⁴, where only approximate distances between parganas were respected. We used some of the Parganas and Sarkars names and their modern correspondences to correct the maps of the Subas (Provinces) of ‘Vizapour’ (Bijapur) and ‘Aiderabad’ (Hyderabad), corresponding to the

RESEARCH COMMUNICATIONS

Table 1. Details of soil organic carbon (SOC) stocks and surface areas per land-cover type at three different dates corresponding to (i) potential; (ii) 18th century situations with two land-cover hypotheses (a) and (b); and (iii) recent situations

	(i) Potential		(iia) 1770		(iib) 1770		(iii) 1977		(iii) 1999	
	Area (sq. km)	SOC stock (PgC)	Area (sq. km)	SOC stock (PgC)	Area (sq. km)	SOC stock (PgC)	Area (sq. km)	SOC stock (PgC)	Area (sq. km)	SOC stock (PgC)
Nilgiris 'sholas'	1,294	0.013	1,294	0.014	1,294	0.014	461	0.005	461	0.005
Nilgiris grasslands	552	0.006	552	0.005	552	0.005	295	0.002	295	0.002
Evergreen forests continuum	46,789	0.287	44,000	0.282	44,000	0.282	3,656	0.023	3,509	0.023
Plateau moist deciduous forests	5,995	0.044	4,800	0.035	4,800	0.035	3,284	0.024	2,955	0.022
Plateau dry deciduous forests	33,748	0.204	5,500	0.036	5,500	0.036	3,068	0.020	2,916	0.019
Secondary/disturbed evergreen forest			5,000	0.032	5,000	0.032	13,250	0.076	11,594	0.073
Plateau cropland (rainfed)			2,520	0.009	5,040	0.018	18,940	0.069	9,700	0.035
Plateau cropland (irrigated)			250	0.001	560	0.003	9,343	0.047	18,524	0.089
Non-forest coastal lowlands, cultivated or not			1,500	0.006	1,500	0.006	5,835	0.025	6,545	0.025
Degraded forests, wastelands			22,943	0.085	20,108	0.091	23,907	0.105	22,121	0.100
Plantations			0	0	0	0	4,639	0.032	6,617	0.045
Miscellaneous/unclassified			–	–	–	–	647	–	643	–
Water	106		125	0.000	130	0.000	1,159	0	3,604	0
Total	88,484	0.554	88,484	0.522	88,484	0.520	88,484	0.429	88,484	0.437

present-day states of Karnataka, Kerala, Tamil Nadu and Goa plus the southern part of Maharashtra. Part of the corrected map corresponding to the Goa–Belgaum region in the north of the study area is given, Figure 2a, along with a modern map, Figure 2b, showing the present-day administrative divisions.

In 1807, after visiting a large part of the study area during 1800–1801, Buchanan²⁵ described a flourishing country, well cultivated, and gave numerous statistics, including those reporting that coffee was being imported in India^{25,26}.

In the late 1820s, the first commercial coffee plantation was established in Shimoga district, Karnataka²⁶, hence initiating the industrial mode of natural resource use in the study area¹⁷. From 1820 onwards, this mode of resource use had an important effect on deforestation in the evergreen ecosystems. Black pepper, coffee and cardamom were grown under the shade of medium-elevation evergreen forests after the undergrowth was cleared, while stands of low and medium-elevation evergreen forests were completely removed for rubber. Tea and cinchona (*Cinchona succirubra*) were grown above 1500 m altitude. Several timber or fuel wood species (*Eucalyptus* spp., *Acacia* spp.) were also planted in clear-felled areas or at the expense of high-altitude grasslands. The development of plantations probably led to compensation between SOC losses in deforested areas and gains in reforested ones as it is observed nowadays¹⁶.

A special mention can be made of deforestation of deciduous forests due to the construction of the Indian railway network during the second part of the 19th century. Quoting different sources, Gadgil and Guha¹⁷ reported that in India 'in the 1870s, it was calculated that

well over a million sleepers were required annually'. According to Legris²⁷ and Meher-Homji²⁸, these requirements led to selective felling of sal (*Shorea robusta*) in the dry deciduous forests of central India and of teak (*Tectona grandis*) in those of South India, and to such an extent that teak completely disappeared from forest stands in some places. In many other places, settled cultivation was still prevailing, with clear felling of forests for the extension of cultivated lands, forest degradation for fire wood and animal grazing, with a net loss of biomass and SOC stock.

From this concise historical scenario of the Indian peninsula, the following assumptions can be made: (i) shifting cultivation and nomadic pastoralism prevailed long before 1770, impacting most of the deciduous formations of the plateau and leading to a great extension of degraded forests and wastelands; (ii) in 1770, the Nilgiris was still the territory of the Badagas, Todas and Kotas communities²⁹, which had limited impact on their environment²⁹; (iii) at that time there were no plantations in the study area; (iv) deforestation for food crops had marginally concerned evergreen forests and (v) irrigation was limited to the eastern side of the study area and covered less than 10% of the plateau cropland (in 1977, the area of irrigated cropland represented 33% of the total area of plateau cropland¹⁶).

The SOC dynamics of the study area can then be outlined at three reference dates. Till 1000 BC, all the area was covered by forests except for high-altitude grasslands; at that time, SOC stock can be considered close to the potential SOC stock, i.e. 0.554 PgC. Estimates of recent stocks gave 0.430 PgC for the 1999 situation¹⁶, a figure stable during 1977–1999 as already mentioned¹⁶.

The third date corresponds to the 1770s situation, i.e. at the time Gentil prepared his maps, and is also within the period that other studies^{5,6} considered as corresponding to potential vegetation. To estimate SOC stocks at this point we simulated a land-cover scenario (Table 1) with plateau-cultivated areas limited to 10% (a) and 20% (b) of their 1999 extension. Both hypotheses gave estimates close to 0.520 PgC in 1770. Therefore about 0.030 PgC emitted before 1770 was neglected in most of the published scenarios^{4–6} and *de facto* added to the more recent emissions. In other words, previous calculations transfer 25% of the total emissions from the pre-1770 period to the recent one. Without minimizing the seriousness of the depletion of carbon stocks in the contemporary period, we argue that the role of the pre-19th century times should be taken into account for an accurate evaluation of the variation in SOC in South India, as also in other countries of Asia with a long history.

- IPCC, *Climate Change 2007: Synthesis Report*, Cambridge University Press, Cambridge, 2007.
- IPCC, Revised 1996 IPCC guidelines for national greenhouse gas inventories, OECD, Paris, 1997.
- IPCC, IPCC guidelines for national greenhouse gas inventories, prepared by the National Greenhouse Gas Inventories Programme, IGES vol. 4, Agriculture, Forestry and other Land Use, Tokyo, 2006.
- IPCC, *Climate Change 2001: The Scientific Basis*, Cambridge University Press, Cambridge, 2001.
- Houghton, R. A. and Hackler, J. L., Emissions of carbon from forestry and land use changes in tropical Asia. *Global. Change Biol.*, 1999, **5**, 481–492.
- Van Minnen, J. G., Goldewijk, K. K., Stehfest, E., Eickhout, B., van Drecht, G. and Leemans, R., The importance of three centuries of land-use change for the global and regional terrestrial carbon cycle. *Climate Change*, 2009, **97**, 123–144.
- Chhabra, A., Palria, S. and Dadhwal, V. K., Soil organic carbon pool in Indian forests. *For. Ecol. Manage.*, 2003, **173**, 187–199.
- Leemans, R., Global datasets collected and compiled by the Biosphere Project, Working Paper, International Institute for Applied Analysis, Laxenburg, Austria, 1990.
- De Fries, R. S., Field, C. B., Fung, I., Collatz, G. J. and Bounoua, L., Combining satellite data and biogeochemical models to estimate global effects of human-induced land-cover change on carbon emissions and primary productivity. *Global. Biogeochem. Cycles*, 1999, **13**, 803–815.
- Pascal, J.-P., Shyam Sunder, S. and Meher-Homji, V. M., Forest map of South India (1 : 250,000 scale), sheets Shimoga and Mercara–Mysore. Institut français de Pondichéry, Puducherry, 1982.
- Pascal, J.-P., Shyam Sunder, S. and Meher-Homji, V. M., Forest map of South India (1 : 250,000 scale), sheet Belgaum–Dharwar–Panaji, Institut français de Pondichéry, Puducherry, 1984.
- Ramesh, B. R., De Franceschi, D. and Pascal, J.-P., Forest map of South India (1 : 250,000 scale), sheet Coimbatore–Thrissur, Institut français de Pondichéry, Puducherry, 2002.
- Pascal, J.-P., Wet evergreen forests of the Western Ghats of India. Ecology, structure, floristic composition and succession. Institut français de Pondichéry, Travaux de la Section Scientifique et Technique, Puducherry, 1988, vol. 20bis.
- Gaussen, H., Legris, P., Labroue, L., Meher-Homji, V. M. and Viart, M., International map of vegetation and of environmental conditions at 1 : 1,000,000 scale, sheet Mysore, explanatory booklet. Institut français de Pondichéry, Hors série 7, Puducherry, 1966.
- Krishnan, P. *et al.*, Organic carbon stock map for South Indian soils: a multifactorial approach. *Curr. Sci.*, 2007, **93**, 706–710.
- Lo Seen, D., Ramesh, B. R., Nair, K. M., Martin, M., Arrouays, D. and Bourgeon, G., Soil carbon stocks, deforestation and land-cover changes in the Western Ghats biodiversity hotspot (India). *Global Change Biol.*, 2010, **16**, 1777–1792.
- Gadgil, M. and Guha, R., *This Fissured Land. An Ecological History of India*, Oxford University Press, New Delhi, 1983.
- Ellis, E. C. and Ramankutty, N., Putting people in the map: anthropogenic biomes of the world. *Front. Ecol. Environ.*, 2008, **6**, 439–447.
- Van Oppen, M., Tank irrigation in southern India: adapting a traditional technology to modern socioeconomic conditions. In *Alfisols in Semi-Arid Tropics: A Consultants' Workshop on the State of the Art and Management Alternative for Optimizing the Productivity of SAT Alfisols and Related Soils* (eds Pathak, P., El-Swaify, S. A. and Singh, S.), ICRISAT, Patancheru, 1987, pp. 89–93.
- Fuller, D. Q., Agricultural origins and frontiers in South Asia: a working synthesis. *J. World Prehist.*, 2006, **20**, 1–86.
- Fuller, D. Q., Dung mounds and domesticators: early cultivation and pastoralism in Karnataka. In *South Asian Archaeology, Vol. I, Prehistory* (eds Jarrige, C. and Lefèvre, V.), Edition Recherche sur les Civilisations, Paris, 2006, pp. 117–127.
- Randhawa, M. S., *A History of Agriculture in India*, Indian Council of Agricultural Research, New Delhi, 1980.
- Sreenivasa Murthy, H. V. and Ramakrishnan, R., *History of Karnataka*, S. Chand & Co Ltd, New Delhi, 1983.
- Gole, S., Maps of Mughal India drawn by Colonel Jean-Baptiste-Joseph Gentil, agent from the French Government to the Court of Shuja-ad-daula at Faizabad, in 1770. Manohar Publications, New Delhi, 1988.
- Buchanan, F., *A Journey through the Countries of Mysore, Canara and Malabar, T. Cadell and W. Davies, London, 1807*, Asian Educational Services, First reprint, New Delhi–Madras, 1988 reprint.
- Government of Karnataka, *Karnataka State Gazetteer of India*, Chikmagalur District, Government Press, Bangalore, 1981.
- Legris, P., La végétation de l'Inde. Ecologie et flore. Travaux de la Section Scientifique et Technique, Institut Français de Pondichéry, Puducherry, 1963, vol. 6.
- Meher-Homji, V. M., History of vegetation of peninsular India. *Man Environ.*, 1989, **13**, 1–10.
- Emeneau, M. B., Linguistics and botany in the Nilgiris. In *Blue Mountains Revisited, Cultural Studies on the Nilgiri Hills* (ed. Hockings, P.), Oxford University Press, New Delhi, 1997, pp. 74–105.

ACKNOWLEDGEMENTS. Part of this work was funded by the Indo-French Centre for Promotion of Advanced Research, New Delhi, in the form of a research project. Among the collaborators of the project, who are not authors of the present paper, we thank P. Krishnan and S. Srinivas (Bangalore Centre of the National Bureau of Soil Survey), D. Arrouays and M. P. Martin (Orléans Centre of the National Institute for Agronomic Research, INRA, France) and F. Landy (University of Paris Ouest-Nanterre-La Défense, France).

Received 13 February 2012; accepted 15 May 2012