

# Role of earthworm in improving soil structure and functioning

D. K. Sharma<sup>1,\*</sup>, S. Tomar<sup>1</sup> and D. Chakraborty<sup>2</sup>

<sup>1</sup>Centre for Environment Science and Climate Resilient Agriculture, ICAR-Indian Agricultural Research Institute, New Delhi 110 012, India

<sup>2</sup>Division of Agricultural Physics, ICAR-Indian Agricultural Research Institute, New Delhi 110 012, India

**Earthworms commonly occur within the soil. They alter physico-chemical and biological regimes of the soil through their activities, such as burrowing, casting, feeding and propagating, and therefore are known as ‘ecosystem engineers’. Through their activities, they provide a number of ecosystem services which are ecologically and socio-economically important. This article reviews the role of earthworms in improving soil structural and functional properties, which serves as key determinants of soil ecosystem services and economic benefits for the farmers. A methodology of economic evaluation of the agro-ecosystem services provided by earthworms has been demonstrated. Further, the information gaps and future research have been discussed for ensuring sustainable agro-ecosystems management.**

**Keywords:** Earthworm, economic evaluation, soil ecosystem services, soil aggregation, sustainable agriculture.

ECOSYSTEM services include an array of direct and indirect benefits provided by natural ecosystems for the well-being of human societies and represent part of the economic value of the planet<sup>1</sup>. Soil-forming processes account for more than one third of the average value of the total ecosystem services (33 Trillion US\$)<sup>2</sup>. In general, biodiversity is the basis of these ecosystem services at every aspect. Earthworms play an important role in the soil-forming process and therefore considered as ‘keystone species’<sup>3</sup> and described as ‘ecosystem engineers’<sup>4</sup>, a vastly popular term proposed only in 1994 (refs 5, 6). They are also called ‘ecosystem services managers’ as they act as a potential partner of human beings<sup>7</sup>.

Earthworms have been categorized into three principal ecological groups depending upon the ecosystem processes and the services they provide (Figure 1). For example, most common epigeic species produce soil roughness and macro-pores by inhabiting litter and producing casts at the soil surface. Anecic species extend down the soil layer as they live in vertical burrows. The third category is the endogeic species, which make horizontal or randomly oriented burrows through all the layers of the upper soil and feed on decayed organic matter<sup>8</sup>.

Earthworms act as a catalyst in enhancing the crop yield (Table 1), but their effect depends on the amount of crop residue, earthworm density and rate of fertilization<sup>9</sup>. However, they always provide materials and energy outputs which have monetary values. They also carry out a range of functions by supporting life through regulating bio-geochemical cycles and other biosphere processes such as decomposition, climate, pollution remediation and biodiversity interaction. Moreover, earthworm’s life processes lead to several other services which are necessary for the functioning and self-sustainability of agro-ecosystems. These services include primary production, nutrient cycling, soil structure development and soil hydraulic functions, to name a few. Earthworm is also an important component of education to students to demonstrate the decomposition and recycling of organic matter. The services of earthworms are still not recognized and there is a lack of awareness of its ecological and economic benefits.

The objectives of this review are (i) to discuss the role of earthworms in transforming soils and (ii) to evaluate the potential role of earthworms in the agro-ecosystem services.


## Earthworm services to the soil and agro-ecosystem

### *Soil structural development*

Development of soil structure from micro- to macro-scale is a function of interacting physical forces of soil water content, actions of large biota like plant roots and earthworms, presence of organic matter in the soil and tillage<sup>10</sup>. The most prominent effect of the earthworm is on the reorganization of soil structure while moving through the soil, feeding on the soil<sup>11</sup> and release of material due to their low assimilation efficiency.

Earthworms can either make the soil compact or loose, depending on the species and its interaction with the soil. Anecic and endogeic species are responsible for a majority of physical improvements in soil structure through cast production in vertical and horizontal burrows. Cast production modifies soil bulk density through incorporation of organic matter into the soil. For example, an

\*For correspondence. (e-mail: dsharma@iari.res.in)



**Figure 1.** Ecosystem service cascade linking biophysical measurements of natural capital with ultimate benefits to society<sup>64</sup>.

endogeic earthworm, *Reginaldia omodeoi* was reported to increase soil bulk density from 1.24 to 1.31 g cm<sup>-3</sup> and from 1.37 to 1.48 g cm<sup>-3</sup> in two different studies<sup>12</sup>. A significant increase was also observed in bulk density from 1.12 to 1.23 g cm<sup>-3</sup>. A resulting decrease in porosity from 58% to 53% in the presence of a tropical endogeic earthworm, *Pontoscolex corethrurus* was also observed<sup>13</sup>. In a long-term field experiment under tropical condition, interaction between soil compacting (*R. omodeoi*) and de-compacting (small *Eudrilidae* family) species helped in maintaining the soil structure<sup>14</sup>. Earthworms also affect size distribution of soil aggregates. De-compacting earthworms (e.g. *Millsonia anomala*) destroyed the macro-aggregates formed by the compacting ones, whereas compacting earthworms acted similarly with the casts of the de-compacting earthworms<sup>15</sup>. This indicates that there exists a wide variability in the action of earthworms in regulating soil structure dynamics. It was reported that compacting earthworms (e.g. *R. omodeoi*), inoculated under yam and maize culture increased the relative proportion of aggregates >2 mm from 30% to 54% and 25% to 42% respectively<sup>16,17</sup>. Earthworms are estimated to produce casts at the rate of about 40–100 tonnes ha<sup>-1</sup> year<sup>-1</sup> and thus, immensely contribute to the formation of stable soil aggregates<sup>18</sup>. They could convert 18–42% of soil material into macro-aggregates only within two months<sup>19</sup>.

It is obvious that earthworms improve soil porosity as well. In a study, fine (<0.4 mm) fraction of soil aggregates was predominant in a treatment with no earthworm population, compared to the treatments where earthworms were present<sup>15</sup>. It was also observed that endogeic earthworms modified soil porosity to control the water flow in the soil, and the effect depended on the equilibrium between cast production and soil degradation<sup>20</sup>.

### Soil water retention

The dwelling activity of earthworms modifies soil aggregates and increases stable macro-pores<sup>21</sup> which improves water infiltration to the soil. Interestingly, each ecologically distinct group of earthworms has different impacts on soil water flow due to the dissimilar burrowing activity of the groups<sup>22</sup>. A study reported enhanced drying of the topsoil by anecic species *Lumbricus terrestris* and endogeic species *Aporrectodea caliginosa*, while the presence of *Lumbricus rubellus* tended to increase the soil water status<sup>23</sup>. In comparison to other species, *A. caliginosa* helps in enhancing the soil infiltration and faster water movement to the subsoil layers due to its high dwelling activity in the soil.

Earthworms interact with plant species composition of agro-ecosystem. A certain plant functional group along with earthworm biomass could significantly alter the spatial and temporal variations in soil hydraulic properties while soil texture had no impact on such properties<sup>24</sup>.

### Soil carbon sequestration

Earthworms induce the primary production of ecosystems by enhancing the nutrient release in the soil, thereby contributing to climate change regulation activities<sup>25</sup>. This action could be due to enrichment of organic matter, nutrients and water in the gut of the earthworm species, which encourage microbial activity<sup>26</sup>. These microbial byproducts facilitate in binding the soil particles into stable aggregates<sup>27</sup>. Thus, earthworms help in maintaining a higher C content through the formation of stable aggregate and stabilizing C by preventing its otherwise rapid degradation and loss as CO<sub>2</sub> and/or CH<sub>4</sub> (ref 28). An

**Table 1.** Role of earthworms in modifying soil physical, chemical and biological properties

Properties	Parameters	Earthworm species involved	Change in the property (% over control)	
Physical	Aggregates	<i>Eudrillus eugeniae</i> <sup>78</sup>	-19.50 (Micro), -76.97 (Macro)	
	Bulk density	<i>Millsonia anomala</i> <sup>79</sup>	71.54	
		<i>M. anomala</i> <sup>79</sup>	8.029	
		<i>Aporrectodea longa</i> , <i>Aporrectodea tuberculata</i> and <i>Bimastos parvus</i> , <i>Phoenicodrilus taste</i> <sup>80</sup>	0.71	
	Hydraulic conductivity	<i>Aporrectodea caliginosa</i> <sup>81</sup>	8.06	
<i>Lumbricus rubellus</i> <sup>81</sup>		5.44		
<i>Octolasion cyaneum</i> <sup>80</sup>		28.54		
Water holding capacity	<i>Megascolex megascolex</i> , <i>Eudrilus eugeniae</i> , <i>Eisenia fetida</i> <sup>77</sup>	61		
Chemical	Soil organic carbon	<i>Pheretima alaxandri</i> <sup>38</sup>	2.66	
		<i>E. eugeniae</i> <sup>78</sup>	27.22	
		<i>Pontoscolex corethrurus</i> <sup>82</sup>	21	
		Exotic species ( <i>A. tuberculata</i> , <i>Lumbricus terrestris</i> ) <sup>83</sup>	34.81	
		<i>A. longa</i> , <i>A. tuberculata</i> and <i>B. parvus</i> , <i>P. taste</i> <sup>80</sup>	64.14	
	pH	<i>M. anomala</i> <sup>79</sup>	-1.36	
		<i>Pheretima alaxandri</i> <sup>38</sup>	10.61	
		NA <sup>84</sup>	27.08	
		<i>A. caliginosa</i> <sup>85</sup>	5.6	
	Nitrogen content	<i>A. longa</i> , <i>A. tuberculata</i> and <i>Bimastos parvus</i> , <i>Phoenicodrilus taste</i> <sup>80</sup>	-4.54	
		<i>E. eugeniae</i> <sup>78</sup>	50	
		<i>P. corethrurus</i> <sup>82</sup>	2.04	
		Exotic species ( <i>A. tuberculata</i> , <i>L. terrestris</i> ) <sup>3</sup>	48.52	
		<i>A. caliginosa</i> <sup>85</sup>	1.9	
	Phosphorus content	<i>A. longa</i> , <i>A. tuberculata</i> and <i>Bimastos parvus</i> , <i>P. taste</i> <sup>80</sup>	47.11	
<i>M. anomala</i> <sup>79</sup>		-1.51		
<i>Metaphire guillelmi</i> <sup>86</sup>		-28.69		
<i>P. alaxandri</i> <sup>38</sup>		26.66		
Potassium content	<i>E. eugeniae</i> <sup>78</sup>	72.47		
	<i>A. longa</i> , <i>A. tuberculata</i> and <i>B. parvus</i> , <i>P. taste</i> <sup>80</sup>	20.12		
Sodium content	<i>M. guillelmi</i> <sup>82</sup>	-27.11		
	<i>P. alaxandri</i> <sup>38</sup>	80.66		
Biological	Microfauna	<i>E. eugeniae</i> <sup>78</sup>	64.07	
		NA	3.64	
		<i>A. longa</i> , <i>A. tuberculata</i> and <i>B. parvus</i> , <i>P. taste</i> <sup>80</sup>	3.64	
	Soil organic matter Enzymatic activities	NA	Bacteria <sup>87</sup>	26.25
			Fungi <sup>87</sup>	17.39
			Actinomycetes <sup>87</sup>	15
		<i>A. caliginosa</i>	Protozoa <sup>88</sup>	-18.24
			Rotifers <sup>88</sup>	-90
			Nematodes <sup>88</sup>	-83.80
		<i>P. corethrurus</i>	Nematodes <sup>89</sup>	-20.60
		NA	Azotobacter <sup>90</sup>	93.16
		NA <sup>6</sup>		26.22
		NA	Cellulase <sup>87</sup>	8.73
			Amylase <sup>87</sup>	57.82
			Invertase <sup>87</sup>	79.87
	Urease <sup>87</sup>	18.67		
	Protease <sup>8</sup>	20		
<i>A. caliginosa</i>	Cellulase <sup>88</sup>	-17.2		
	Dehydrogenase <sup>88</sup>	79.26		
	Protease <sup>88</sup>	98.27		

**Table 2.** Role of earthworms in CO<sub>2</sub> and N<sub>2</sub>O emission from the soil

Continents	Country	Emission (mg kg <sup>-1</sup> )	
		CO <sub>2</sub>	N <sub>2</sub> O
Asia	China <sup>27</sup>	1079.8	0.4227
America	Canada <sup>65</sup>	340.0	0.105
	Canada <sup>66</sup>	2568.0	0.818
	Mexico <sup>67</sup>	11.3	2.600
Europe	Netherlands <sup>68</sup>	881.0	0.056–0.789
	Netherlands <sup>69</sup>	473.9–593.6	1.397
	Netherlands <sup>70</sup>	302.0	0.973
Africa	Antsirabe, Madagascar <sup>71</sup>	48.0	3.800
Oceania	Australia <sup>72</sup>	1510.6	1.055

increase in soil organic C from 16.1 to 17.9 g C kg<sup>-1</sup> and from 12.4 to 14.7 g C kg<sup>-1</sup> after the introduction of *Lumbricus terrestris* to a chisel-tilled soil under maize-soya bean rotation has been reported<sup>29</sup>.

Increase in C emission from fields with a good earthworm population was also widely reported (Table 2). A meta-analysis indicated that although earthworms sequester C in macro-aggregates, it also increases CO<sub>2</sub> emissions by 37% through aerobic respiration<sup>30</sup>. A decline in organic matter from an arable land with high earthworm population was recorded after 15 years<sup>31</sup>. It appears that earthworms can stabilize the soil organic matter when organic residues are regularly added to the soil<sup>32</sup>. Addition of organic matter is recommended rather than the inoculation of earthworms into the soil alone, to improve the soil fertility in the agro-ecosystem<sup>33</sup>. A majority of Indian soils are low in organic C status, and therefore it is likely that earthworm services can be realized only when organic inputs were added to the soil to encourage soil C sequestration.

### Nutrient cycling in soil

A number of studies highlighted the role of earthworms in nitrogen flow in the agro-ecosystem by increasing the nitrous oxide (N<sub>2</sub>O) emissions (Table 2). The flux of N by earthworms was estimated at 63 kg ha<sup>-1</sup> year<sup>-1</sup> in no-tillage system<sup>34</sup>, and 10 and 41 kg ha<sup>-1</sup> year<sup>-1</sup> in monoculture and organically fertilized agro-ecosystems<sup>35,36</sup>.

Earthworms exhibit profound influence on soil microbial community and increase nitrogen mineralization in soil<sup>37</sup>. However, earthworm effects on N-cycling is determined by cropping system and fertilizer types, and by the predominant species and species interactions in the soil<sup>38</sup>. Therefore the earthworm species and substrate present in the soil largely determine the N-immobilization and/or N-mineralization<sup>39</sup>. Hence, agroecosystem management becomes extremely important<sup>40</sup>. Mineralization of organic substances is assisted through ingestion, fragmentation and transportation of the partially decomposed

plant residues by earthworms which increase the availability of nutrients in the soil<sup>41</sup>. Plant growth is improved through increase in nutrient composition in plant tissues<sup>42</sup>, and through higher nutrient release in the soil and uptake<sup>43</sup>. Microbial activity is increased in the presence of earthworms, leading to nitrogen-mineralization and immobilization through direct or indirect benefits by earthworms<sup>44</sup>.

Earthworms were also reported to increase the availability of phosphorus by facilitating changes in the biogeochemical status of P in burrow-linings<sup>45</sup>. In a pot experiment with earthworms, higher levels of plant available P resulted in increased plant growth from 15.7 to 16.8 g dry biomass per pot<sup>46</sup>.

### Soil biodiversity interactions

Earthworms are an integral part of the soil ecosystem, and therefore, exhibit intimate functional interactions with microbes. The effect is manifested through three basic mechanisms: comminuting, burrowing and casting, and grazing and dispersal, which change the soil's physico-chemical and biological regimes. Earthworms possibly condition the substrate which promotes microbial activities.

Earthworms induce the nematode population in the soil and nematode trophic structure<sup>47</sup> which help in assimilating amino acids with other compounds<sup>48</sup>. In contrast, *Lampito mauritii* decreased the numbers of nematodes<sup>49</sup>. Micro-arthropod population in soils e.g. *Collembola* in soils was increased in the presence of earthworms<sup>50</sup>. Micro-arthropods get attracted by earthworms through their action on surrounding soil, where water availability, aeration and pores size are improved<sup>51</sup>.

### Pollution remediation

Earthworms are reported to bio-accumulate high concentrations of heavy metals like Cd, Hg, Pb, Cu, Mn, Ca, Fe

**Table 3.** Heavy metal concentration (mg kg<sup>-1</sup>) in different species of earthworms

Species	Cd	Cu	Pb	Zn
<i>Pheretimaposthuma</i> <sup>73</sup>	0.0498	6.092	0.0173	1.92
<i>Pheretima californica</i> <sup>74</sup>	9.18	74.68	670.55	657.75
<i>Aporrectodea caliginosa</i> <sup>75</sup>	11.6–102.9	17.9–35.9	1.9–182.8	556–3381
<i>Lumbricus rubellus</i> <sup>75</sup>	7.7–26.3	16.0–37.6	0.5–37.9	667.9–2645
<i>Aporrectodea tuberculata</i> <sup>76</sup>	–	361	–	631
<i>Eisenia foetida</i> <sup>76</sup>	–	249	–	439
<i>Lumbricus terrestris</i> <sup>38,77</sup>	–	39.0	31.0	132
	–	62.0	10110	1550

and Zn in their tissues<sup>52</sup> by ingesting them with soil. This converts ionic state of these metals and make them available for plants after earthworm excretion (Table 3). In a study, higher amounts of heavy metals were found in earthworms collected from the roadside and mining sites<sup>53</sup>. Earthworms accumulate higher amounts of Zn and Cd<sup>54</sup> and reduce the concentrations of Cr, Cu, Zn and Pb to the limits set by USEPA in 60 days using vermicompost sludge (biosolids)<sup>55</sup>. Thus earthworms may be regarded as bio-indicators for evaluation of soil health.

#### Production enhancement

Earthworm's role in plant growth has been widely demonstrated<sup>43</sup>, possibly due to increased nutrient availability in soils. A meta-analysis revealed the role of earthworms in increasing crop yield by 25% and above ground biomass by 23% (ref. 9). Leaf weight, cob biomass and the number of cobs of maize crop also increased by 40%, 152% and 130% respectively, in the presence of *Millsonia anomala*; increased leaf production of rice was recorded by 14% in the presence of *Chuniodrilu zizelae*. Association of *M. anomala* and *C. zizelae* had better impact than their individual effect<sup>56</sup>.

#### Considerations for economic quantification

Diversity, abundance and biomass are the major parameters to be considered for quantifying the role of earthworms in ecosystem services. We have so far discussed the ecosystem services by earthworms. Here we develop a methodology for quantification and valuation of these services in monetary terms. The methodology covers following five major steps.

##### Identifying agroecosystem services

Although direct benefits of earthworms are disguised, their contribution to the ecological modification of agroecosystem and to the economy of farmers is well-recognized. Identification of major benefits is the prime step for quantification and valuation of ecosystem services, considering the socio-economic and ecological measures of an individual service.

##### Developing markets

The next step is to develop the market for identified services. In general, it is crucial to ensure the proximity of the service in terms of land management approach, farmers' intervention (fertilizer and irrigation inputs), human activity, demography and transfer to market cost for generating a service market<sup>57</sup>. Six steps have been identified for creating ecosystem services market: (1) generation of demand; (2) definition of unit and supply; (3) definition and establishment of payment mechanism; (4) establishment of supporting institutions; (5) feedback and environment<sup>58</sup> and (6) stability and conditionality of programme implementation<sup>59</sup>.

##### Assigning monetary values

Stakeholders may assign values to the ecosystem services based on the markets and the discharge of ecological and socioeconomic services. Many studies have given values to a particular service by measuring the earthworm's population in subsoil or topsoil. Earthworm population was used to estimate the quantity of fertile soil annually<sup>60</sup> depending on biomass and soil turnover estimates<sup>61,62</sup> respectively. Besides, at an average earthworm biomass of 75 g m<sup>-1</sup>, grassland productivity was approximated to US\$ 152 million in 2.25 m ha of land<sup>63</sup>.

##### Consideration of external drivers

Once monetary values are assigned, the external drivers (climate change, land-use pattern, etc.) affecting the soil processes may be considered for cost-benefit analysis. These factors influence the soil processes leading to lesser natural capital stocks.

##### Trade-off analysis

This is a process of quantifying the relationship between key indicators associated with changes in agro-ecosystems and relevant policies. A meta-analysis indicated that earthworms increase CO<sub>2</sub> emissions rather than its

sequestration<sup>30</sup>. We must take note of these issues which affect policy-making, and influence the stakeholders or decision makers. A trade-off analysis to quantify the indicator's effect on agro-ecosystem and in policy-making is therefore and must be considered.

## Conclusions

We need larger and more robust data on soil biodiversity to understand the ecosystem services by soil fauna, i.e. microorganisms and micro-arthropods. The earthworm is an essential component in sustainable agricultural systems, although major gaps remain in understanding their role in providing ecosystem services. Therefore, there is a need to identify all the ecosystem services provided by each ecological group of earthworms along with other soil biota. The existing and new knowledge on earthworms and related ecosystem services need to be integrated. Additionally, the farm-level management practices should be enhanced to promote ecosystem services<sup>64</sup>. Cost-benefit or trade-off analysis must be accounted. It is also necessary to update the knowledge of monetary values generated by earthworm population at local or regional scale for economic evaluation of ecosystem services.

1. MA (Millennium Ecosystem Assessment), *Ecosystems and Human Well-being: Synthesis*, Island Press, Washington DC, 2005.
2. Costanza, R. *et al.*, The value of world's ecosystem services and natural capital. *Nature*, 1997, **387**, 253–260.
3. Lyons, K. G., Brigham, C. A., Traut, B. H. and Schwartz, M. W., Rare species and ecosystem functioning. *Conserv. Biol.*, 2005, **19**(4), 1019–1024.
4. Hale, C. M., Frellich, L. E., Reich, P. B. and Pastor, J., Effects of European earthworm invasion on soil characteristics in northern hardwood forest of Minnesota, USA. *Ecosystem*, 2008, **8**, 911–927.
5. Jones, C. G., Lawton, J. H. and Shachak, M., Organisms as ecosystem engineers. *Oikos*, 1994, **69**, 373–386.
6. Wright, J. P. and Jones, C. G., The concept of organisms as ecosystem engineers ten years on: progress, limitations and challenges. *BioSci.*, 2006, **56**, 203–209.
7. Byers, J. E. *et al.*, Using ecosystem engineers to restore ecological systems. *Trends Ecol. Evol.*, 2006, **21**, 493–500.
8. Lee, K. E., *Earthworms their Ecology and Relationships with Soils and Land Use*, Academic Press, Sydney, 1985.
9. van Groenigen, J. W., Lubbers, I. M., Vos, H. M. J., Brown, G. G., De Deyn, G. B. and van Groenigen, K. J., Earthworms increase plant production: a meta-analysis, 2014; doi:10.1038/srep06365.
10. Milleret, R., Le Bayon, R. C. and Gobat, J. M., Root, mycorrhiza and earthworm interactions: their effects on soil structuring processes, plant and soil nutrient concentration and plant biomass. *Plant Soil*, 2009, **316**, 1–12.
11. Martin, A. and Marinissen, J. C. Y., Biological and physico-chemical processes in excrements of soil animals. *Geoderma*, 1993, **56**, 331–347.
12. Lavelle, P. *et al.*, Effects of earthworms on soil organic matter and nutrient dynamics at a landscape scale over decades. In *Earthworm Ecology* (ed. Edwards, C. A.). CRC Press, Boca Raton, 2004.
13. Alegre, J., Pashanasi, B. and Lavelle, P., Dynamics of soil physical properties in Amazonian agroecosystems inoculated with earthworms. *Soil Sci. Soc. Am. J.*, 1996, **60**, 1522–1529.
14. Blanchart, E. *et al.*, Effects of earthworms on soil structure and physical properties. In: *Earthworm Management in Tropical Agroecosystems* (eds Lavelle, P., Brussaard, L. and Hendrix, P.), CAB International, Wallingford, 1999, pp. 149–172.
15. Blanchart, E., Lavelle, P., Braudeau, E., Le Bissonnais, Y. and Valentin, C., Regulation of soil structure by geophagous earthworm activities in humid savannas of Côte d'Ivoire. *Soil Biol. Biochem.*, 1997, **29**, 431–439.
16. Gilot, C., Effects of a tropical geophagous earthworm, *Millsonia anomala* (Megascolecidae), on soil characteristics and production of a yam crop in Ivory Coast. *Soil Biol. Biochem.*, 1997, **29**, 353–359.
17. Gilot-Villeneuve, C., Lavelle, P. and Ganry, F., Effects of a tropical geophagous earthworm, *Millsonia anomala*, on some soil characteristics, on maize-residue decomposition and on maize production in Ivory Coast. *Appl. Soil Ecol.*, 1996, **4**, 201–211.
18. Pietz, R. I., Peterson, J. R., Prater, J. E. and Zenz, D. R., Metal concentrations in earthworms from sewage sludge-amended soils at a strip mine reclamation site. *J. Environ. Qual.*, 1984, **13**(4), 651–654.
19. Zangerle, A., Pando, A. and Lavelle, P., Do earthworms and roots cooperate to build soil macroaggregates? A microcosm experiment. *Geoderma*, 2011, **167–168**, 303–309.
20. Bottinelli, N., Henry-des-Tureaux, T., Hallaire, V., Mathieu, J., Benard, Y., Duc Tran, T. and Jouquet, P., Earthworms accelerate soil porosity turnover under watering conditions. *Geoderma*, 2010, **156**, 43–47.
21. Bastardie, F., Capowiez, Y., de Dreuzy, J.-R. and Cluzeau, D., X-ray tomographic and hydraulic characterization of burrowing by three earthworm species in repacked soil cores. *Appl. Soil Ecol.*, 2003, **24**, 3–16.
22. Edwards, W. M., Shipitalo, M. J., Owens, L. B. and Norton, L. D., Effect of *Lumbricus terrestris* L. burrows on hydrology of continuous no-till corn fields. *Geoderma*, 1990, **46**, 73–84.
23. Ernst, G., Felten, D., Vohland, M. and Emmerling, C., Impact of ecologically different earthworm species on soil water characteristics. *Eur. J. Soil Biol.*, 2009, **45**, 207–213.
24. Fischer, C. *et al.*, How do earthworms, soil texture and plant composition affect infiltration along an experimental plant diversity gradient in grassland? *PLoS ONE*, 2014, **9**(6): e98987.
25. Amosse, J., Turberg, P., Kohler-Milleret, R., Gobat, J.-M. and Le Bayon, R.-C., Effects of endogeic earthworms on the soil organic matter dynamics and the soil structure in urban and alluvial soil materials. *Geoderma*, 2015, **243–244**, 55–57.
26. Haynes, R. J. and Fraser, P. M., A comparison of aggregate stability and biological activity in earthworm casts and uningested soil as affected by amendment with wheat or lucerne straw. *Eur. J. Soil Sci.*, 1998, **49**, 629–636.
27. Tianxiang, L., Huixin, L., Tong, W. and Feng, H., Influence of nematodes and earthworms on the emissions of soil trace gases (CO<sub>2</sub>, N<sub>2</sub>O). *Acta Ecol. Sin.*, 2008, **28**, 993–999.
28. Leon, Y. S., Lugo-Perez, J., Wise, D. H., Jastrow, J. D. and Gonzalez-Meler, M. A., Aggregate formation and carbon sequestration by earthworms in soil from a temperate forest exposed to elevated atmospheric CO<sub>2</sub>: A microcosm experiment. *Soil Biol. Biochem.*, 2014, **68**, 223–230.
29. Shuster, W. D., Subler, S. and McCoy, E. L., Deep burrowing earthworm additions changed the distribution of soil organic carbon in a chisel-tilled soil. *Soil Biol. Biochem.*, 2001, **33**, 983–996.
30. Lubbers, I. M., van Groenigen, K. J., Fonte, S. J., Six, J., Brussaard, L. and van Groenigen, J. W., Greenhouse-gas emissions from soils increased by earthworms. *Nature Clim. Change*, 2013, **3**, 187–194.

31. Riley, H., Pommeresche, R., Eltun, R., Hansen, S. and Korsaeht, A., Soil structure, organic matter and earthworm activity in a comparison of cropping systems with contrasting tillage, rotations, fertilizer levels and manure use. *Agric. Ecosyst. Environ.*, 2008, **124**, 275–284.
32. Fonte, S. J. and Six, J., Earthworms and litter management contributions to ecosystem services in a tropical agroforestry system. *Ecol. Appl.*, 2010, **20**(4), 1061–1073.
33. Blouin, M. *et al.*, A review of earthworm impact on soil function and ecosystem services. *Eur. J. Soil Sci.*, 2013, **64**(2), 161–182.
34. Parmelee, R. W. and Crossley Jr, D. A., Earthworm production and role in the nitrogen cycle of a no-tillage agroecosystem on the Georgia piedmont. *Pedobiologia*, 1988, **32**, 353–361.
35. Rosswall, T. and Paustian, K., Cycling of nitrogen in modern agricultural systems. *Plant Soil*, 1984, **76**, 3–21.
36. Whalen, J. K., Parmelee, R. W. and Subler, S., Quantification of nitrogen excretion rates for three lumbricid earthworms using <sup>15</sup>N. *Biol. Fertil. Soils*, 2000, **32**, 347–352.
37. Barois, I., Lavelle, P. and Kajondo, J. K., Adaptive strategies and short term effects of selected earthworm species, selection of particles. In Conservation of Soil Fertility in Low-Input Agricultural Systems of the Humid Tropics by Manipulating Earthworm Communities (ed. Lavelle, P.), IRD, Bondy, France, 1992, Report of the CCE Project No. TS2.0292-F(EDB), pp. 35–67.
38. Bhadauria, T. and Saxena, K. P., Role of earthworms in soil fertility maintenance through the production of biogenic structures. *Appl. Environ. Soil Sci.*, 2010, p. 7; doi:10.1155/2010/816073.
39. Brown, G. G. *et al.*, Effects of earthworms on plant production in the tropics. In Earthworm Management in Tropical Agro Ecosystems (eds Lavelle, P., Brussaard, L. and Hendrix, P.), CABI Publishing, Wallingford, UK, 1999, pp. 87–147.
40. Fonte, S. J., Kong, A. Y. Y., van Kessel, C., Hendrix, P. F. and Six, J., Influence of earthworm activity on aggregate-associated carbon and nitrogen dynamics differs with agroecosystem management. *Soil Biol. Biochem.*, 2007, **39**, 1014–1022.
41. Sharpley, A. N. and Syers, J. K., Potential role of earthworm casts for the phosphorus enrichment of run-off waters. *Soil. Biol. Biochem.*, 1976, **8**, 341–346.
42. Stefan S., Effects of earthworms on plant growth: patterns and perspectives. The 7th International Symposium on Earthworm Ecology, Cardiff, Wales, Submitted 6 September 2002.
43. Mayilswami, S. and Reid, B., Effect of earthworms on nutrients dynamics in soil and growth of crop. *Res. J. Chem. Environ.*, 2010, **14**(2), 39–44.
44. Blair, J. M., Parmelee, R. W. and Lavelle, P., Influences of earthworms on biogeochemistry. In *Earthworm Ecology and Biogeography in North America* (ed. Hendrix, P. F.), CRC Press, Boca Raton, 1995, pp. 127–158.
45. Le-Bayon, R. C. and Binet, F., Earthworms change the distribution and availability of phosphorus in organic substrates. *Soil Biol. Biochem.*, 2006, **38**, 235–246.
46. Vos, H. M. J., Ros, M. B. H., Koopmans, G. F. and van Groenigen, J. W., Do earthworms affect phosphorus availability to grass? A pot experiment. *Soil Biol. Biochem.*, 2014, **79**, 34–42.
47. Domínguez, J., Parmelee, R. W. and Edwards, C. A., Interactions between *Eisenia andrei* (Oligochaeta) and nematode populations during vermicomposting. *Pedobiologia*, 2003, **47**, 53–60.
48. Pokarzhevskii, A. D., Zaboev, D. P., Gordienko, S. A. and Ganin, G. N., Are earthworms ecosystemivorous? *Soil Biol. Biochem.*, 1997, **29**, 559–567.
49. Dash, M. C., Senapati, B. K. and Mishra, C. C., Nematode feeding by tropical earthworms. *Oikos*, 1980, **34**, 322–325.
50. Satchell, J. E., Lumbricidae. In *Soil Biology* (eds Burges, A. and Raw, F.), Academic Press, New York, NY, 1967, pp. 259–322.
51. Devliegher, W. and Verstraete, W., Microorganisms and soil physico-chemical conditions in the drilosphere of *Lumbricus terrestris*. *Soil Biol. Biochem.*, 1997, **29**, 1721–1729.
52. Ireland, M. P., Heavy metals uptake in earthworms; earthworm ecology. Chapman & Hall, London, 1983.
53. Sinha, R. K., Bharambe, G. and Chaudhari, U., Sewage treatment by vermin-filtration with synchronous treatment of sludge by earthworms: a low cost sustainable technology over conventional systems with potential for decentralization. *Environmentalist*, 2008, **28**, 409–420.
54. Hartenstein, R., Neuhauser, E. F. and Collier, J., Accumulation of heavy metals in the earthworm *E. foetida*. *J. Environ. Qual.*, 1980, **9**, 23–26.
55. Contreras-Ramos, S. M., Escamilla-Silva, E. M. and Dendooven, L., Vermicomposting of biosolids with cow manure and oat straw. *Biol. Fertil. Soils*, 2005, **41**, 190–198.
56. Derouard, L., Tondoh, J., Vilcosqui, L. and Lavelle, P., Effects of earthworm introduction on soil processes and plant growth. *Soil Biol. Biochem.*, 1997, **29**(3/4), 541–545.
57. Boyd, J. and Wainger, L., Measuring Ecosystem Service Benefits: The Use of Landscape Analysis to Evaluate Environmental Trades and Compensation, Discussion. Paper 02-63, Resources for the Future, Washington DC, 2003.
58. Landell-Mills, N. and Porras, I., Silver Bullet or Fools' Gold? International Institute for Environment and Development, London, 2002.
59. Wunder, S., Payments for Environmental Services: Some Nuts and Bolts. CIFOR Occasional Paper No. 42, Center for International Forestry Research, Jakarta, 2005.
60. Sandhu, H. S., Wratten, S. D., Cullen, R. and Case, B., The future of farming: the value of ecosystem services in conventional and organic arable land. An experimental approach. *Ecol. Econ.*, 2008, **64**, 835–848.
61. Fraser, P. M., Williams, P. H. and Haynes, R. J., Earthworm species, population size and biomass under different cropping systems across the Canterbury Plains, New Zealand. *Appl. Soil Ecol.*, 1996, **3**, 49–57.
62. Pimentel, D. *et al.*, Environmental and economic costs of soil erosion and conservation benefits. *Science*, 1995, **267**(5201), 1117–1123.
63. Boag, B. and Neilso, R., Impact of New Zealand flatworm on agriculture and wildlife in Scotland. *Proc. Crop Prot. Northern Br.*, 2006, pp. 51–55.
64. Keith, A. M. and Robinson, D. A., Earthworms as natural capital: ecosystem service providers in agricultural soils. *Economology J.*, 2012, **2**, 91–99.
65. Speratti, A. B. and Whalen, J. K., Carbon dioxide and nitrous oxide fluxes from soil as influenced by anecic and endogeic earthworms. *Appl. Soil Ecol.*, 2008, **38**, 27–33.
66. Speratti, A., Whalen, J. and Rochette, P., Earthworm influence on carbon dioxide and nitrous oxide fluxes from an unfertilized corn agroecosystem. *Biol. Fert. Soils*, 2007, **44**, 405–409.
67. Contreras-Ramos, S. M., Alvarez-Bernal, D., Montes-Molina, J. A., van Cleemput, O. and Dendooven, L., Emission of nitrous oxide from hydrocarbon contaminated soil amended with waste water sludge and earthworms. *Appl. Soil Ecol.*, 2009, **41**, 69–76.
68. Rizhiya, E., Bertora, C., van Vilet, P. C. J., Kuikman, P. J., Faber, J. H. and van Groenigen, J. W., Earthworm activity as a determinant for N<sub>2</sub>O emission from crop residue. *Soil Biol. Biochem.*, 2007, **39**, 2058–2069.
69. Giannopoulos, G., Pulleman, M. M. and van Groenigen, J. W., Interactions between residue placement and earthworm ecological strategy affect aggregate turnover and N<sub>2</sub>O dynamics in agricultural soil. *Soil Biol. Biochem.*, 2010, **42**, 618–625.
70. Bertora, C., van Vliet, P. C. J., Hummelink, E. W. J. and van Groenigen, J. W., Do earthworms increase N<sub>2</sub>O emissions in ploughed grassland? *Soil Biol. Biochem.*, 2007, **39**, 632–640.
71. Chapuis-Lardy, L. *et al.*, Effect of the endogeic earthworm *Pontoscolex corethrurus* on the microbial structure and activity related

- to CO<sub>2</sub> and N<sub>2</sub>O fluxes from a tropical soil (Madagascar). *Appl. Soil Ecol.*, 2010, **45**, 201–208.
72. Chan, Y. C., Sinha, R. K. and Wang, W., Emission of greenhouse gases from home aerobic composting, anaerobic digestion and vermicomposting of household wastes in Brisbane (Australia). *Waste Manage. Res.*, 2011, **29**(5), 540–548.
  73. More, B. C. and Patole, S. S., Bioaccumulation of heavy metals through earthworms from water hyacinth (*Eichhornia crassipes* Solm) contaminated vermicompost. *J. Environ. Res. Develop.*, 2012, **7**(2A), 1047–1050.
  74. Zhen-zhong, W., You-mei, Z., Yong-can, G., Wei-sheng, X. and Zhong-wu, L., Monitoring of soil heavy metals by earthworm. *J. Environ. Sci.*, 1998, **10**(4), 437–444.
  75. Dai, J., Becquer, T., Rouiller, J. H., Reversat, G., Bernhard-Reversat, F., Nahmani, J. and Lavelle, P., Heavy metal accumulation by two earthworm species and its relationship to total and DTPA-extractable metals in soils. *Soil Biol. Biochem.*, 2004, **36**, 91–98.
  76. Lukkari, T., Aatsinki, M., Väisänen, A. and Haimi, J., Toxicity of copper and zinc assessed with three different earthworm tests. *Appl. Soil Ecol.*, 2005, **30**, 133–146.
  77. Munnoli, P. M. and Bhosle, S., Water-holding capacity of earthworms' vermicompost made of sugar industry waste (press mud) in mono- and polyculture vermireactors. *Environmentalist*, 2011, **31**(4), 394–400.
  78. Kruapakdee, A. W., Iwai, C. B. and Ta-Oun, M., The influence of earthworm and rice stubble on soil aggregate, carbon sequestration and soil fertility in sandy loam paddy soil in Northeast Thailand. 4th ICERD-International Conference on Environmental and Rural Development, 2013.
  79. Lavelle, P. *et al.*, Large-scale effects of earthworms on soil organic matter and nutrient dynamics. In *Earthworm Ecology* (eds Edwards, C. A.), St. Lucie Press, 2010, pp. 103–122.
  80. Castellanos-Navarrete, A., Rodriguez-Aragones, C., de Goede, R. G. M., Kooistra, M. J., Sayre, K. D., Brussaard, L. and Pulleman, M. M., Earthworm activity and soil structural changes under conservation agriculture in central Mexico. *Soil Tillage Res.*, 2012, **123**, 61–70.
  81. Francis, G. S. and Fraser, P. M., The effects of three earthworm species on soil macroporosity and hydraulic conductivity. *Appl. Soil Ecol.*, 1998, **10**(1–2), 11–19.
  82. Zhang, B., Liu, Y., Li, C., Hu, F. and Velde, B., Long-term fertilization influences on clay mineral composition and ammonium adsorption in a rice paddy soil. *Soil Sci. Soc. Am. J.*, 2010, **74**, 1239–1247.
  83. Wironen, M. and Moore, T. R., Exotic earthworm invasion increases soil carbon and nitrogen in an old-growth forest in southern Quebec. *Can. J. For. Res.*, 2006, **36**, 845–854.
  84. Avizinin, E. J., The indirect effect of earthworms on methane flux in two temperate pasture soils. Master's thesis, University of Rhode Island, 2012.
  85. McColl, H. P., Hart, P. B. S. and Cook, F. J., Influence of earthworms on some soil chemical and physical properties, and the growth of ryegrass on a soil after topsoil stripping—a pot experiment. *New Zeal. J. Agric. Res.*, 2012, **25**(2), 239–243.
  86. Zhang, B. G., Li, G. T., Shen, T. S., Wang, J. K. and Sun, Z., Changes in microbial biomass C, N and P and enzyme activities in soil incubated with the earthworms *Metaphire guillelmi* or *Eisenia fetida*. *Soil Biol. Biochem.*, 2000, **32**, 2055–2062.
  87. Devi, S. H., Vijayalakshmi, K., Jyotsna, K. P., Shaheen, S. K., Jyothi, K. and Rani, M. S., Comparative assessment in enzyme activities and microbial populations during normal and vermicomposting. *J. Environ. Biol.*, 2009, **30**(6), 1013–1017.
  88. Aira, M., Monroy, F. and Dominguez, J., Effects of two species of earthworms (*Allolobophora* sp.) on soil systems: a microfaunal and biochemical analysis. *Pedobiologia*, 2003, **47**, 877–881.
  89. Villenave, C., Rabary, B., Kichenin, E., Djigal, D. and Blanchart, E., Earthworms and plant residues modify nematodes in tropical cropping soils (Madagascar): a Mesocosm experiment. *Appl. Environ. Soil Sci.*, 2010, **7**.
  90. Suhane, R. K. Vermicompost (In Hindi); Pub. of Rajendra Agricultural University, Pusa, Bihar, 2007, p. 88.

ACKNOWLEDGEMENTS. Our thanks to the Department of Science and Technology, New Delhi for providing financial support for this work. We also thank the Indian Agricultural Research Institute, New Delhi, India for providing the platform to conduct the study.

Received 29 June 2016; revised accepted 10 June 2017

doi: 10.18520/cs/v113/i06/1064-1071