and sub-acute apex. Apiscopic and basiscopic margins straight, upper basal angle-free and rounded, lower basal angle decurrent. Midrib is distinct, running from base to apex.

According to Sahm<sup>4</sup> E. plana is the commonest and one of the most characteristic fossil of the Raghavapuram and Superamatur Shales (Kota stage). It is also reported from Bansa in Madhya Pradesh (Jabalpur Stage). Baksi<sup>5</sup> reported it from Raghavapuram mudstones in the West Godavari District of Andhra Pradesh. Mahabale and Satyanarayana<sup>6</sup> reported it from Raghudevapuram in East Godavari District. This is the first report from Uppugunduru in Prakasam District. It appears that E. plana has wider distribution in the Jurassic beds of Andhra Pradesh.

Specimen No. -UPG/50/86

Locality -Uppugunduru, Prakasam

District (A.P.)

Horizon -Upper Jurassic.

## 7 February 1987

- 1. Vagyani, B., Curr. Sci., 1985, 54, 14.
- 2. Halle, T. G., Ergebnisse, 1913, 3, 14.
- 3. Seward, A.C., Fossil Plants, 1919.
- 4. Sahni, B., Palaeontologia indica, 1928, N.S. XI.
- 5. Baksi, S. K., Palaeobotanist, 1969, 16, 3.
- 6. Mahabale, T. S. and Satyanarayana, T., Geophytology, 1979, 9, 1.

## CYTOGENETICS OF A PENTAPLOID HYBRID OF SOLANUM SCABRUM MILL. AND S. VILLOSUM MILL.

A. GANAPATHI and G. R. RAO

Department of Botany, Bharathidasan University,
Tiruchirapalli 620 024, India.

SOLANUM SCABRUM Mill. the 'garden huckleberry' is a tribasic (2n=6x=72) species of the S. nigrum complex. Its origin and interrelationship with other species of the section Solanum are not well known<sup>1</sup>. The present note reports the crossability between S. scabrum and S. villosum (2n=4x=48).

The plants of S. scabrum were raised from seeds supplied by Dr Z. O. Gbile, Forest Research Institute, Ibadan, Nigeria, and a stock of S. villosum was raised from local populations. Crosses between the two species were successful only when S. scabrum, the higher chromosomal form, was used as a maternal parent. A total number of 100 cross

pollinations yielded 5 fruits with 2 seeds, but only one germinated and grew to maturity. The  $F_1$  hybrid was a sterile pentaploid with 2n=5x=60 chromosomes. Pollen fertility was as low as 4.7%. The hybrid exhibited a wide range of meiotic irregularities. At diakinesis and metaphase I, besides bivalents, univalents and multivalents were recorded in more than 90% of the cells. The mean association of chromosomes, per cell, at metaphase I, was  $10.12_I + 23.81_{II} + 0.70_{III} + 0.04_{IV}$ . Anaphase I and the subsequent stages of meiosis were irregular and characterized by laggards, chromatin bridges and unequal distribution of chromosomes at poles. The parents showed normal meiosis and bivalent associations.

The breakdown of meiosis and high pollen sterility of the hybrid seems to be due to chromosomal cause, because the parental species differ in chromosome number<sup>2</sup>. However, the vigorous hybrid indicates existence of genetic harmony between the parental forms. The occurrence of about 23 bivalents in a majority of the pollen mother cells indicates the homology between 23 chromosomes each of S. villosum and S. scabrum, thereby demonstrating the participation of S. villosum genome in ancestry of S. scabrum.

The participation of S. villosum genome in the origin of S. nigrum is already established<sup>1,3,4</sup>. It is of interest to note that S. scabrum and S. nigrum were readily crossable with each other producing fertile hybrids<sup>4</sup>. Apparently S. nigrum and S. scabrum have originated from the same or identical ancestors; changes responsible for morphological divergence accumulated during the course of evolution by structural changes of chromosomes as well as gene mutations.

## 21 February 1987; Revised 12 May 1987

- 1. Edmonds, J. M., The biology and taxonomy of Solanaceae, Linn. Soc. Symp. series, No. 7, 1979, p. 529.
- 2. Stebbins, G. L., Chromosomal evolution in higher plants, Edward Arnold Ltd., London, 1971, p. 216.
- 3. Ganapathi, A. and Rao, G. R., Curr. Sci., 1985, 54, 1242.
- 4. Ganapathi, A., Ph.D. thesis, University of Madras, Madras, 1984.