

14. Lensky, Y., *Comp. Biochem. Physiol.*, 1971, **B50**, 335.
15. Pentz, S. and Kling, H., *J. Insect Physiol.*, 1972, **18**, 2277.
16. deLoof, A. and deWilde, J., *J. Insect Physiol.*, 1970, **16**, 157.
17. Brown, J. J. and Chippendale, G. M., *Insect Biochem.*, 1978, **8**, 359.
18. Kunkel, J. G., *Comp. Biochem. Physiol.*, 1975, **B51**, 177.
19. Kunkel, J. G. and Lawler, D. M., *Comp. Biochem. Physiol.*, 1974, **B47**, 697.
20. McCormick, F. W. and Scott, A., *Experientia*, 1966, **22**, 228.

ON THE IDENTITY OF A LYMANTRIID DEFOLIATOR OF CASHEW AND COCOA IN SOUTH INDIA

G. RAMASESHIAH and RAMESH BALI
CAB International Institute of Biological Control,
Post Bag 2484, Hebbal Agricultural Farm Post,
Bangalore 560 024, India.

AMONG several species of the genus *Lymantria* Hubner occurring in India, *Lymantria obfuscata* Walker, defoliating alder, apple, apricot, false acacia, oak, pear, peach, plum, poplar, walnut, willow and other fruit trees in Jammu and Kashmir, and in Himachal Pradesh, is considered as a highly destructive pest comparable with *Lymantria dispar* L. in Europe and America. Several parasites of it are known in India¹⁻⁷. Records of its occurrence on cashew and cocoa³⁻⁵ in Tamil Nadu and Karnataka interested the authors while looking for a multivoltine brood of this species for studies on inter-specific competition of parasites of *L. obfuscata* available in Kashmir and Himachal Pradesh. An intensive survey for *L. obfuscata* in Tamil Nadu and Karnataka was made in 1985-86.

Large numbers of a destructive lymantriid larvae were collected on cashew in the plantations of Narimanam, Samakottai and Vriddhachalam (Tamil Nadu), cashew and cocoa in plantations around Vittal (Karnataka) and reared in the laboratory. Both male and female moths and the parasites obtained were referred to the CIE for determination. The moths were identified as *Lymantria ampla* Walker. The other lymantriid pests on cocoa recorded by Premkumar and Radhakrishnan Nair⁵ were *Euproctis subnotata* Wlk., *E. guttata* Wlk. and *Dasychira mendosa* HB. Figures of a larva, a gravid female and a male moth of *L. obfuscata* were

provided by them. These figures agree with *L. ampla* which is also reported as a pest of cocoa from Sri Lanka².

Neonatal larvae of *L. obfuscata* from Kashmir failed to develop on cashew leaves in the laboratory. The male and female genitalia of *L. obfuscata* and of the moth from cashew in the south differ significantly. The braconids *Aleiodes* sp., *Apanteles obliquae* Wilkinson, *Apanteles* sp. (*glomeratus* group), a eulophid *Euplectrus* sp., a chalcid *Brachymeria porthetrialis* Joseph, Narendran & Joy and the tachinids *Blepharipa* sp., *Carcelia* sp., *Exorista* sp. and *Palexorista* sp. were reared from *L. ampla* collected on cashew in Tamil Nadu. The parasite complex of *L. obfuscata* is different with the exception of *A. obliquae*. Around Bangalore, braconids *Apanteles* sp., *Apanteles* sp. (Gr.A) and *Meteorus* sp., tachinids *Blepharella lateralis* Macq., *Blepharipoda zebina* Walker, *Carcelia* sp.,? *buitenzorgiensis* Baranov, *Carelia* sp.,? *C. buitenzorgiensis* Baranov, *Drino (Prosturmia)* sp., *D (P.) lucagus* Walker, *Exorista japonica* Townsend, *E. sp.*,? *larvarum* L. and the ichneumonids *Bari-chneumon* sp., *Enicospilus* sp., *Pimpla poesia* Cameron, the chalcids *Brachymeria banksi* (Ashmead), *B. deesensis* Cameron, *B. euploae* Westwood and a eulophid *Trichospilus* sp. parasitised *L. ampla* on *Ficus religiosa* and *Casuarina equisetifolia*. In the light of these observations the record of *L. obfuscata* on cashew and cocoa in the south is obviously erroneous. The damage potential of *L. ampla* to cashew and cocoa warrants consideration of biological control.

This research has been financed in part by a grant from the US Department of Agriculture under Cooperative Agricultural Research Grant Program (PL 480).

Thanks are due to Drs. K. M. Harris, J. D. Holloway, T. Huddleston and A. K. Walker of the CAB International Institute of Entomology for identifying the specimens.

7 May 1987

1. Dharmadhikari, P. R., Ramaseshiah, G. and Achan, P. D., *Entomophaga*, 1985, **30**, 398.
2. Entwistle, P. F., *Pests of cocoa*, Longmans, London, 1972, p. 779.
3. Misra, M. P. and Basu Choudhuri, J. C., *Indian For.*, 1974, **100**, 391.
4. Premkumar, T., *Curr. Sci.*, 1974, **43**, 395.
5. Premkumar, T. and Radhakrishnan Nair, C. P., CPCRI Reg. Stn. Vittal, *Publn. No.* 174, 1978, 3.

6. Rishi, N. D. and Shah, K. A., *J. Entomol. Res.*, 1985, 9, 82.
 7. Zutshi, M. F., *J. Bombay Nat. Hist. Soc.*, 1966, 64, 126.

RECORD OF MEALYBUG SPECIES ON GRAPEVINE IN KARNATAKA*

M. MANI and T. S. THONTADARYA†

Division of Entomology and Nematology, Indian Institute of Horticultural Research, Hessaraghatta Lake Post, Bangalore 560 089, India.

† No. 256, 1st 'N' Block, Rajajinagar, Bangalore 560 010, India.

In recent years, the mealybugs have caused severe damage to the grapevine *Vitis vinifera* in certain parts of India^{1,2}. They are found on the leaves, shoots, nodes, bunches, under loose bark of grapevine etc. A survey was carried out in the major grape-growing areas of Karnataka to determine the species of mealybugs infesting grapevine during January–March in 1984–1986.

The present investigation revealed the presence of four species of mealybugs.

1. *Planococcus citri* (Risso): It was found to cause more than 75% damage to the bunches of the variety Black Champa at Hessaraghatta. In the nursery, severely affected vines died due to this mealybug attack. This is the first record on grapes in India though reports about this as a major pest of grapevine in more than ten countries are available.

2. *Maconellicoccus hirsutus* (Green): It was commonly occurring on grape bunches in all the major grape-growing areas of Karnataka causing 80% damage. This species was reported to occur on grapevine in Andhra Pradesh¹, Karnataka² and New Delhi³.

3. *Nipaecoccus viridis* (Newstead): It was observed in two vineyards around Bangalore. A maximum of 5% damage due to this mealybug was observed. This mealybug species has been reported earlier in New Delhi⁴.

4. *Dysmicoccus brevipes* (Ckll.): It was of minor importance around Bangalore. However it has not been reported earlier as a pest of grapevine elsewhere.

The authors are grateful to Dr W. J. Williams, Commonwealth Institute of Entomology, London for identifying the mealybug species.

22 June 1987

1. Azam, K. M., *Indian J. Entomol. (special issue)*, 1983, 2, 387.
 2. Manjunath, T. M., *FAO Plant Prot. Bull.*, 1985, 32, 74.
 3. Fletcher, T. B., *Sci. Rep. Agric. Res. Inst.*, 1919, p. 86.
 4. Subba Rao, B. R., Sangwar, H. S., Abbasi, O. A., Singh, V. and Ksheer Sagar, A. M., *Indian J. Entomol.*, 1965, 27, 109.

RECORDS OF BLEPHARELLA LATERALIS MACQUART AND CARCELIA SP. — TWO INDIGENOUS PARASITIDS OF SPILOSOMA (=DIACRISIA) OBLIQUA WALKER FROM BIHAR (INDIA)

NAWIN KUMAR and R. P. YADAV

Department of Entomology, Rajendra Agricultural University, Dholi-Campus, Muzaffarpur 843 121, India.

BIHAR hairy caterpillar, *Spilosoma* (= *Diacrisia*) *obliqua* Walker is a polyphagous pest which is known to cause serious damages to a variety of economically important crops in India. With a view to exploring the possibility of its biocontrol, preliminary investigations were initiated to record its naturally occurring parasitoids. Pest larvae of different age-groups were collected from the field at weekly intervals and reared in the laboratory on the natural foods.

Of the two tachinid parasitoids, viz, *Carcelia* sp. and *Blepharella lateralis* Macquart, the former was recorded as larval parasitoid while the latter as larval-pupal one. Both these parasitoids were found active from mid-August to end-November, with the combined rate of natural parasitization varying from 3 to 6% only. Usually one and occasionally two fully grown maggots of *Carcelia* sp. came out from the body of one parasitized pest larva of pre-pupal stage through its lateral region and underwent pupation within 6–12 hr. On the other hand, the pest larvae parasitized by *B. lateralis* pupated normally but usually 1 or 2 days earlier than the healthy ones. But the pupae of parasitized caterpillars later became blackish and the parasitoid fly emerged usually one from one such pupa. Occurrence of *Carcelia* sp. as the larval parasitoid of *S. obliqua* was earlier reported from Punjab¹ and Jabalpur² while *B. lateralis* was recorded for the first time from India.

* Contribution No. 82/87.