

**GENUS *PSEUDENDACLONIUM* WILLE
(CHAETOPHORALES) FROM INDIA**

R. K. TIWARI*, H. M. SHUKLA,
U. C. PANDEY and G. L. TIWARI**

Department of Botany, Bareilly College,
Bareilly 243 001, India.


* Tara Silk Development Project, Bahraich 271 801, India.

** Department of Botany, University of Allahabad, Allahabad 211 002, India.

THE genus *Pseudendoclonium* established by Wille¹ represents a strongly reduced and adapted relative

of *Stigeoclonium* Kuetz. and *Endoclonium* Szymanski. *Pseudendoclonium* is known to have only four species^{2,3} with a marine form viz. *P. submarinum* Wille^{1,4} and to date this genus is not reported from India. The present paper deals with two new records of species for Indian flora viz. *P. akinetum* Tupa and *P. prostratum* Tupa.

During our investigation of freshwater epiphytic algal forms of Allahabad and its nearby localities, the authors came across two species of the genus viz. *P. akinetum* Tupa and *P. prostratum* Tupa which are hitherto not reported from India. Irregular tufts of



Figures 1-8. 1. *Pseudendoclonium akinetum* Tupa showing quadriflagellate zoospore; 2. Young thalli; 3. Mature cells; 4. Zoosporangia having zoospores; 5. Akinetes; 6. *Pseudendoclonium prostratum* Tupa showing quadriflagellate zoospore; 7. Young thallus, and 8. Mature thallus.

P. akinetum and *P. prostratum* were found growing epiphytically in knee-deep water of the temporary ponds of Jhusi (Allahabad) and Mouri (Pratapgarh) Districts respectively.

P. akinetum Tupa: Thallus tufts forming, mainly consists of prostrate filaments of usually 6–10 cells but occasionally up to 20 cells in a single filament. Prostrate filaments are irregularly branched and may produce extremely short erect filaments of usually 3–4 cells. Often young thalli have a distinct main creeping filament which may have branches of 3–4 elongated and terminal tapering cells (figure 2). Each of the elongated young cells may be up to 5 μ wide and 10 μ long and becomes subspherical and enlarged rapidly up to 14 μ in diameter (figure 3). All cells possess a single dark green parietal band-shaped chloroplast which covers the entire surface of the cell embedded in which there is a single prominent centrally placed pyrenoid.

Asexual reproduction occurs by zoospores as well as akinetes. Zoosporangia are spherical, usually enlarged up to 17 μ in diameter (figure 4). From each zoosporangium, 4 to 8 zoospores are released. Zoospores are quadriflagellate measuring up to 5 μ in diameter and 8 μ in length. They are pear-shaped and have a cup-shaped chloroplast with a posterior pyrenoid and a distinct anteriorly placed eye spot (figure 1). Akinetes are formed by enlargement of cells and accumulation of starch granules. The cell wall gets thickened and becomes characteristically rust brown in colour (figure 5). Akinetes show unipolar germination.

Epiphytic on blades of aquatic grasses.

P. prostratum Tupa: Thallus is much prominent as compared to *P. akinetum* Tupa. Prostrate system consists of richly branched pseudoparenchymatous single-layered disc. Mature thalli possess numerous erect branches and thus become cushion-shaped (figures 7 and 8). Cells are 5–10 μ in diameter and 9–15 μ in length. The single parietal chloroplast in each cell contains a single pyrenoid. Central cells of the prostrate system appear angular and rounded due to mutual compression.

Asexual reproduction occurs by zoospores. Zoosporangia are spherical and up to 15 μ diameter. Each zoosporangium produces up to 8 zoospores. Zoospores are quadriflagellate ovoid and have a cup-shaped chloroplast with a single posterior pyrenoid and an eye-spot (figure 6). The zoospores are 5 μ in diameter and 9 μ in length. Epiphytic on *Vallisneria spiralis* Linn.

The authors are grateful to Prof. D. C. Pandey for encouragement and to UGC, New Delhi for financial assistance to one of us (RKT).

4 April 1987

1. Wille, N., *Nor. Vidensk. Skrift I Math-nat.*, 1900, 6, 29.
2. Tupa, D. D., *Nova Hedwigia*, 1974, 46, 60.
3. Printz, H., *Hydrobiologia*, 1964, 24, 274.
4. Visher, W., *Bull. Soz. Bot. Geneva*, 1926, 18, 24.

SPONTANEOUS ALLOTETRAPLOID OF *COIX GIGANTEA* \times *C. AQUATICA*

A. B. SAPRE, A. S. NAIK and S. S. BARVE
Department of Botany, Marathwada University,
Aurangabad 431 004, India.

HYBRIDIZATION followed by chromosome doubling has played a significant role in plant evolution since many allopolyploids with perfectly paired chromosomes have established themselves as fertile and genetically constant new species. Majority of the spontaneous polyploids have now been known to be either allopolyploids or segmental allopolyploids rather than autopolyploids¹⁻³. The true autopolyploids are therefore probably the only ones that are produced artificially—the colchicoids. A spontaneous allotetraploid ($2n = 28$), isolated from the hybrid progeny of *Coix gigantea* ($2n = 18$) \times *C. aquatica* ($2n = 10$), is reported here.

Coix is an oriental relative of the cultivated maize. Three species of *Coix* are reported from India⁴ and two of these, *C. gigantea* ($2n = 20$) from Purandhar⁵ and *C. aquatica* ($2n = 10$) from Mhaismal⁶, are involved in the present study. Nullisomics of *C. gigantea* ($2n = 2$, $2n = 18$) were obtained in large number⁷ through nondisjunction⁸ in a domesticated population maintained at the Botanical Garden of this university. When the aneuploids of *C. gigantea*⁹ were grown in close association with *C. aquatica*, spontaneous interspecific hybrids with varied chromosome numbers were obtained through open pollination¹⁰. The interspecific hybrids were semifertile and when hybrid derivatives were screened cytologically through acetocarmine squashes, individuals with chromosome numbers in the range of $2n = 10$ to $2n = 21$ were isolated¹¹. The present allotetraploid plant was detected among the seed progeny of this population.