

plants sprayed with sterile water served as control. Detached leaves and potted plants were then placed under high humidity conditions. The symptoms appeared on detached leaves and potted plants two and three days after inoculation respectively. The leaves and plants sprayed with sterile water remained healthy. Reisolation from such fungal lesions yielded the same fungus, i.e. *C. gloeosporioides*. The temperature range during pathogenicity tests was 19–28°C.

Although *C. gloeosporioides* has been reported on various aromatic plants²⁻⁷ from different parts of the world, no precise information exists on its occurrence on *P. graveolens*. Therefore, this report constitutes a new record.

The authors thank Dr B. C. Sutton of Commonwealth Mycological Institute, Surrey, UK, for confirming the identity of the pathogen.

15 December 1987

1. Narayana, M. R., Prakasa Rao, E. V. S., Rajeswara Rao, B. R. and Sastry, K. P., *Papai J.*, 1986, 8, 25.
2. Albuquerque, F. C. and Conduru, J. M. P., *Fitotecnia*, 1971, 2, 149.
3. Janardhanan, K. K. and Husain, A., *Curr. Sci.*, 1966, 35, 394.
4. Karunakaran, P. and Nair, M. C., *Plant Dis.*, 1980, 64, 220.
5. Khanna, K. K. and Chandra, S., *Indian J. Mycol. Plant Pathol.*, 1977, 7, 176.
6. Mitidieri, I. Z. N., *Rev. Innesl. Agropecurias*, 1973, 5, 99.
7. Rao, V. G. and Anahosur, K. H., *Indian J. Mycol. Plant Pathol.*, 1971, 1, 79.

RECORD OF A NEW HYMENOPTERAN PARASITOID OF *TRICHOLYGA BOMBYCIS* BECK. (DIPTERA: TACHINIDAE)

G. VEERANNA and H. K. JYOTHI


Karnataka State Sericulture Development Institute,
Thalughattapura, Bangalore 560 062, India

TRICHOLYGA BOMBYCIS Beck, commonly known as uzifly, a major pest of silkworm, *Bombyx mori* prefers to oviposit on III, IV and V instar larvae, killing them and causing damage to the extent of 15–20% to the silk industry¹⁻⁴. Biological methods of insect pest suppression have of late gained increasing acceptance and popularity, as these methods

are specific to the target insect, nonhazardous, more effective and safe. It has been reported that *Nesolynx thymus*, *Trichopria* sp. and *Exoristobia philippinensis* are the pupal parasites of *T. bombycis*⁵⁻⁷. In this report, we record a new hyperparasite, *Dirhinus anthracia* Walker (Chalcididae) of *T. bombycis*.

The pupae of *T. bombycis* were cultured in the laboratory and kept in enamel trays in a single layer for their development. The flies of *D. anthracia* from the field infested some of the uzifly pupae. Flies of *D. anthracia* emerged from these pupae were collected in 250 ml glass beakers covered with muslin cloth and fed on honey and 5% sucrose soaked on cotton pads put on the top of the covered beakers. These flies were then reared at room temperature for further studies. The temperature and humidity recorded in the laboratory ranged from 24 to 31°C and 57 to 92% respectively.

Preliminary observations made on the *D. anthracia* revealed that the female flies are bigger than the males with a broad abdomen and pointed end. The males have narrow abdomen with blunt end (figures 1 and 2). The female mates with the male only once in her lifetime. The pre-mating behaviour showed that the male follows female with wagging movement for about 5 min and mates with her. The duration of mating was 2–5 min, after which the female fly alights on the uzifly pupae, slightly bends her abdomen and pierces her needle-like ovipositor into the pupal case at the intersegmental region to deposit her eggs. Single female lays about 82–157 eggs and lays only one egg per pupa. It was observed that a female fly infested only 1–7 day old pupae and can infest at an average of 105 pupae in her lifetime. The duration of egg stage ranged from 2 to 3 days. The duration of larval and pupal stages ranged from 10 to 13 days and 10 to 12 days respectively. The life-span of the adult flies ranged from 20 to 40 days. The emergence of *D. anthracia* from uzifly pupae started from 20 to 25 days after infestation. The proportion of males and females of *D. anthracia* emerging from the uzifly pupae was 16.4 and 86. Arrhenotokous parthenogenesis was also observed in this species. On parasitization, the larva of *D. anthracia* feed within the host pupa and develop to attain full growth. The fly emerges from the infested pupae by breaking the pupal case. The gravid females of *D. anthracia* were provided with uzi maggots and pupae of different age groups and pupae and larvae of *B. mori* of different age for infestation. The flies emerged only from 1 to 7 day old uzi pupae.



Figures 1 and 2. 1. Adult male fly of *Dirhinus anthracia*, and 2. Adult female fly of *Dirhinus anthracia*.

Investigations on various aspects of *D. anthracia* are in progress to exploit it as a biological control agent to minimize the uzifly population.

The authors are grateful to Dr K. M. Harris, Chief Taxonomist, Commonwealth Institute of Entomology, London, for identification of hyperparasite, *Dirhinus anthracia* Walker.

18 September 1987

1. Mukherjee, N. G., *Handbook of sericulture*, Bengal Secretariat Book Dept., Calcutta, 1919, p. 296.
2. Gosh, E. C., *Silk production and weaving in India*, CSIR Monograph, Delhi, 1949, p. 61.
3. Jolly, M. S., *Uzifly: Its identification, prevention and control*, Central Sericultural Research and Training Institute, Mysore, 1981, p. 1.
4. Kasturi Bai, A. R., Mahadevappa, D., Nirmala, M. R. and Jyothi, H. K., *Curr. Sci.*, 1986, 55, 1038.
5. Pradip Kumar, Venkata Reddy, V., Samson, M. V. and Jolly, M. S., *Curr. Sci.*, 1986, 55, 1040.
6. Veeranna, G., Nirmala, M. R. and Mahadevappa, D., *Curr. Sci.*, 1987, 56, 1031.
7. Veeranna, G., Nirmala, M. R. and Mahadevappa, D., *Sericologia*, 1988, 28, 227.

VIBRIO CHOLERAE 01 AND NON-01 IN SHRIMP "VEIN"—A PRELIMINARY STUDY TO ASSESS ANY POSSIBLE INCIDENCE

P. R. G. VARMA, T. S. G. IYER,
M. AGNES JOSEPH* and SHAJY ZACHARIA*

Central Institute of Fisheries Technology,
Cochin 682 029, India.

* Project on *V. cholerae*, Clo CIFT, Cochin 682 029, India.

THE isolation of *Vibrio cholerae* from shrimps and the consequent rejection of a few consignments by Japan have significantly improved the awareness of the seafood industry about this pathogen. Several queries have been raised about the incidence, source and viability of this organism in marine products.

Since the majority of the consignments rejected by the Japan Quarantine Authorities happened to be headless shrimps, there was a suspicion among some processors that the vein present in the headless shrimps harbours *V. cholerae*. Since no data were available on this aspect, studies were carried out at the Central Institute of Fisheries Technology, Cochin, and the result of the preliminary study is discussed in this paper.

Shrimps harvested from filtration ponds (Chemeenkettu) situated at Parur (Cochin) were used. Twenty-five grams of veins collected from about 100 headless shrimps (*P. indicus* or *P. monodon* of size grade 31/35, 36/40 and 41/50 per lb) constituted one sample.