

HEAVY METAL POLLUTION IN GOMTI RIVER SEDIMENTS AROUND LUCKNOW, UTTAR PRADESH

S. KUMAR

Department of Geology, Lucknow University, Lucknow 226 007, India.

ABSTRACT

The clay fraction ($<2\ \mu\text{m}$) of Gomti river sediments shows relatively higher values for Cu, Mn, Pb, Cr and PO_4 in comparison to background values. Fe, Co and Ni do not show any increase and Cd is not within detectable limits. PO_4 shows very good positive correlation with Cu, Pb, Zn and Mn. It is concluded that municipal waste discharged into the river through drains is responsible for the higher values of heavy metals and PO_4 , but in general Gomti river sediments can be termed as unpolluted with heavy metals.

INTRODUCTION

GOMTI is an alluvial plain river which originates from a swampy area in Pilibhit district and meets Ganga river in Ghazipur district, Uttar Pradesh. Two cities, Lucknow and Jaunpur, are situated on the banks of this river. Lucknow is a big city, with a population of more than 1.5 million and some industrial units. The entire Lucknow municipal waste, including industrial and sewage wastes, flows into the Gomti through small open drains or nalas. This waste alters the quality of the water, which in turn affects the associated sediments of the river. All fine-grained materials with large surface area are capable of accumulating heavy metal ions¹. Thus the clay fraction (i.e. $<2\ \mu\text{m}$ fraction) in a fluvial system is most affected by heavy metal pollution and may acquire a level that may adversely affect the food chain of the aquatic ecosystem. Some of the heavy metals in sediments, specially Pb, Hg, Cu and Cd, are potentially hazardous not only to the aquatic ecosystem but also for human use. In the light of these facts the present paper deals with heavy metal concentration in Gomti river sediments around Lucknow.

The Gomti sediments² are made up of fine sand and mud fractions. The sand is litharenite to lithic graywacke in composition with characteristic heavy minerals indicating metamorphic provenance. Both plagioclases and alkali feldspars are present in moderate amounts in sand and silt fractions. Clay fraction (i.e. $<2\ \mu\text{m}$) shows illite as the main mineral along with kaolinite, montmorillonite and meta-stable chlorite in minor amounts.

SAMPLING

A stretch of 9 km of Gomti river was selected for the present study (figure 1). Eight bank mud samples were collected in March 1987. The locations from where the samples were taken are shown in figure 1.

METHODOLOGY

The $<2\ \mu\text{m}$ fraction was separated from the mud in sedimentation cylinders using double-distilled water at 24°C . This fraction was then dried at 60°C in a porcelain dish. The dried sample (1 g) was digested in 10 ml of 65% pure conc. HNO_3 and




Figure 1. Location map of the samples of Gomti river sediments around Lucknow.




Figure 2. Variation of heavy metal concentration along the Gomti river stretch.

detectable limits. In comparison with world shale standard³ the Gomti clay fraction shows comparable values for Mn, Ni and Cr and low values for Cd and Fe. For Cu, Pb, Zn and Co the Gomti clay fraction has less than thrice the world standard shale values (table 1); however, in comparison with the background values (in the present case values for sample no. 1 are taken as the background values),

these are less than double. Only one sample (sample no. 4) shows PO₄ values more than thrice the background value.

Thus there is a definite increase in heavy metals and PO₄ in Gomti river sediments due to anthropogenic activity but in comparison with background values and world shale standard³ these sediments can be said to be unpolluted with heavy metals. A more than three-fold increase in PO₄, with which the heavy metals show good positive correlation, can be linked to organic pollution due to municipal waste discharged into the river. It is suggested that attempts should be made to minimize this by treating the municipal waste before it is dumped into Gomti river.

ACKNOWLEDGEMENTS

The author is grateful to Prof. S. K. Singh and Prof. G. Müller for encouragement and useful discussions. Thanks are also extended to Messrs M. Gästner, M. Rehauer, L. Haamann and Frau Kuhn for help during the investigation. A fellowship to the author by the Alexander von Humboldt Foundation, W. Germany, to work in the Institut für Sedimentforschung, Heidelberg University, is gratefully acknowledged.

25 January 1988; Revised 12 May 1988

1. Föstner, U. and Whittmann, G. T. W., *Metal pollution in aquatic environment*, Springer-Verlag, Berlin, 1979, p. 486.
2. Kumar, S. and Singh, I. B., *Senckenbergiana marit*, 1979, 10, 145.
3. Turekian, K. K. and Wedepohl, K. H., *Bull. Geol. Soc. Am.*, 1961, 72, 175.

ANNOUNCEMENT

PEPTIDES AS NOVEL PHARMACEUTICALS

(12 and 13 July 1989, Royal Society of Medicine, London, UK)

This meeting will review the current knowledge of peptides in both physiology and pathophysiology and discusses the therapeutic implications for drugs affecting peptide synthesis or action. For details

contact: Dr Renata Duke, IBC Technical Services Ltd, Bath House (3rd Floor), 56 Holborn Viaduct, London EC1A 2EX, UK.