

SUPPLEMENT TO "CURRENT SCIENCE".

Vol. VII]

September 1938

[No. 3

BRITISH ASSOCIATION FOR THE ADVANCEMENT OF SCIENCE.

Cambridge, 1938.

THE PRESIDENTIAL ADDRESS.

Part I.

VISION IN NATURE AND VISION AIDED BY SCIENCE.

Part II.

SCIENCE AND WARFARE.

BY THE RT. HON. LORD RAYLEIGH, Sc.D., LL.D., F.R.S.,

President of the Association.

I.

VISION, AND ITS ARTIFICIAL AIDS AND SUBSTITUTES.

THE last occasion that the British Association met at Cambridge was in 1904, under the presidency of my revered relative, Lord Balfour, who at the time actually held the position of Prime Minister. That a Prime Minister should find it possible to undertake this additional burthen brings home to us how much the pace has quickened in national activities, and I may add, anxieties, between that time and this.

Lord Balfour in his introductory remarks recalled the large share which Cambridge had had in the development of physics from the time of Newton down to that of J. J. Thomson and the scientific school centred in the Cavendish Laboratory, 'whose physical speculations,' he said, 'bid fair to render the closing year of the old century and the opening ones of the new as notable as the greatest which have preceded them.' It is a great pleasure to me, as I am sure it is to all of you, that my old master is with us here to-night, as he was on that occasion. I can say in his presence that the lapse of time has not failed to justify Lord Balfour's words. What was then an intelligent anticipation is now an historical fact.

I wish I could proceed on an equally cheerful note. The reputation of the scientific school in the Cavendish Laboratory has

been more than sustained in the interval under the leadership of one whose friendly presence we all miss to-night. The death of Ernest Rutherford leaves a blank which we can never hope to see entirely filled in our day. We know that the whole scientific world joins with us in mourning his loss.

Lord Balfour's address was devoted to topics which had long been of profound interest to him. He was one of the first to compare the world picture drawn by science and the world picture drawn by the crude application of the senses, and he emphasised the contrast between them. A quotation from his address will serve as an appropriate text to introduce the point of view which I wish to develop this evening.

'So far,' he said, 'as natural science can tell us, every quality or sense or intellect which does not help us to fight, to eat, and to bring up our children, is but a by-product of the qualities which do. Our organs of sense perception were not given us for purposes of research....either because too direct a vision of physical reality was a hindrance, not a help in the struggle for existence....or because with so imperfect a material as living tissue no better result could be attained.'

Some of those who learn the results of modern science from a standpoint of general or philosophical interest come away, I believe, with the impression that what the senses tell us about the external world is shown to be

altogether misleading. They learn, for example, that the apparent or space-filling quality of the objects called solid or liquid is a delusion, and that the volume of space occupied is held to be very small compared with that which remains vacant in between. This is in such violent contrast with what direct observation seems to show that they believe they are asked to give up the general position that what we learn from our senses must be our main guide in studying the nature of things.

Now this is in complete contrast with the standpoint of the experimental philosopher. He knows very well that in his work he does and must trust in the last resort almost entirely to what can be seen, and that his knowledge of the external world is based upon it: and I do not think that even the metaphysician claims that we can learn much in any other way. It is true that the conclusions of modern science seem at first sight to be very far removed from what our senses tell us. But on the whole the tendency of progress is to bring the more remote conclusions within the province of direct observation, even when at first sight they appeared to be hopelessly beyond it.

For example, at the time of Lord Balfour's address some who were regarded as leaders of scientific thought still urged that the conception of atoms was not to be taken literally. We now count the atoms by direct methods. We see the electrometer needle give a kick and we say, 'There goes an atom.' Or we see the path of an individual atom marked out by a cloud track and we see where it was abruptly bent by a violent collision with another atom.

Again, the theory of radioactive decomposition put forward by Rutherford, however cogent it may have seemed and did seem to those who were well acquainted with the evidence, was originally based on indirect inferences about quantities of matter far too small to be weighed on the most delicate balance. Chemists were naturally inclined to feel some reserve; but in due course the theory led to a conclusion which could be tested by methods in which they had confidence—the conclusion, namely, that lead contained in old uranium minerals ought to have a lower atomic weight than ordinary lead and in all probability to be lighter, and on trying this out it proved to be so. More recently we have the discovery of heavy hydrogen with twice the density of ordinary

hydrogen and heavy water which is the source of it.

Lastly, the conclusion that ordinary matter is not really space-filling has been illustrated by the discovery that certain stars have a density which is a fabulous multiple of the density of terrestrial matter. Although this is in some sense a deduction as distinguished from an observation, yet the steps required in the deduction are elementary ones entirely within the domain of the older physics.

This and many other points of view have seemed at first sight to contradict the direct indication of our senses. But it was not really so. They were obtained and could only be obtained by sense indications rightly interpreted. As in the passage from Lord Balfour already quoted the senses were not primarily developed for purposes of research, and we have in large measure to adapt them to that purpose by the use of artificial auxiliaries. The result of doing so is often to reveal a world which to the unaided senses seems paradoxical.

I have chosen for the main subject of this address a survey of some of the ways in which such adaptations have been made. I shall naturally try to interest you by dwelling most on aspects of the subject that have some novelty; but apart from these there is much to be gleaned of historical interest, and when tempted I shall not hesitate to digress a little from methods and say something about results.

I shall begin with a glance at the mechanism of the human eye, so far as it is understood. I shall show how the compromise and balance between different competing considerations which is seen in its design can be artificially modified for special purposes. All engineering designs are a matter of compromise. You cannot have everything. The unassisted eye has a field of view extending nearly over a hemisphere. It gives an indication very quickly and allows comparatively rapid changes to be followed. It responds best to the wave-lengths actually most abundant in daylight or moonlight. This combination of qualities is ideal for what we believe to be nature's primary purpose, that is, for finding subsistence under primitive conditions and for fighting the battle of life against natural enemies. But by sacrificing some of these qualities, and in particular the large field of view, we can enhance others for purposes of research. We

may modify the lens system by artificial additions over a wide range for examining the very distant or the very small. We can supplement and enormously enhance the power of colour discrimination which nature has given us. By abandoning the use of the retina and substituting the photographic plate as an artificial retina, we can increase very largely the range of spectrum which can be utilised. This last extension has its special possibilities, particularly in the direction of using waves smaller than ordinary, even down to those which are associated with a moving electron. By using the photoelectric cell as another substitute for the retina with electric wire instead of optic nerve and a recording galvanometer instead of the brain we can make the impressions metrical and can record them on paper. We can count photons and other particulate forms of energy as well. We can explore the structure of atoms, examine the disintegration of radioactive bodies, and trace out the mutual relation of the elements. Indeed, by elaborating this train of thought a little further almost the whole range of observational science could be covered. But within the compass of an hour or so one must not be too ambitious. It is not my purpose to stray very far from what might, by a slight stretch of language, fall under the heading of extending the powers of the eye.

Most people who have a smattering of science now know the comparison of the eye with the camera obscura, or better, with the modern photographic camera—with its lens, iris, diaphragm, focussing adjustment and ground glass screen, the latter corresponding to the retina. The comparison does not go very far, for it does not enter upon how the message is conveyed to the brain and apprehended by the mind; or even upon the minor mystery of how colours are discriminated. Nevertheless, it would be a great mistake to suppose that the knowledge which is embodied in this comparison was easily arrived at. For example, many acute minds in antiquity thought that light originated in the eye rather than in the object viewed. Euclid in his optics perhaps used this as a mathematical fiction practically equivalent to the modern one of reversing the course of a ray, but other authors appealed to the apparent glow of animal eyes by lamplight, which shows that they took the theory quite literally. The Arabian author Alhazen had more correct ideas and he gave

an anatomical description of the eye, but apparently regarded what we call the crystalline lens as the light-sensitive organ. Kepler was the first to take the modern view of the eye.

The detailed structure of the retina, and its connection with the optic nerve, has required the highest skill of histologists in interpreting difficult and uncertain indications. The light-sensitive elements are of two kinds, the rods and cones. The rods seem to be the only ones used in night vision, and do not distinguish colours. The cones are most important in the centre of the field of view, where vision is most acute, and it seems to be fairly certain that in the foveal region each cone has its own individual nervous communication with the brain. On the other hand, there is not anything like room in the cross-section of the optic nerve to allow us to assign a different nerve fibre to each of the millions of rods. A single fibre probably has to serve 200 of them.

The nervous impulse is believed to travel in the optic nerve as in any other nerve, but what happens to it when it arrives at the brain is a question for the investigators of a future generation.

The use of lenses is one of the greatest scientific discoveries: we do not know who made it. Indeed, the more closely we inquire into this question the vaguer it becomes. Spectacle lenses as we know them are a mediæval invention, dating from about A.D. 1280. Whether they originated from some isolated thinker and experimentalist of the type of Roger Bacon, or whether they were developed by the ingenuity of urban carfts-men, can hardly be considered certain. There are several ways in which the suggestion might have arisen, but a glass bulb filled with water is the most likely. Indeed, considering that such bulbs were undoubtedly used as burning glasses in the ancient world, and that the use of them for reading small and difficult lettering is explicitly mentioned by Seneca, it seems rather strange that the next step was not taken in antiquity. Apparently the explanation is that the magnification was attributed to the nature of the water rather than to its shape. At all events, it may readily be verified that a 4- or 5-inch glass flask full of water, though not very convenient to handle, will give a long-sighted newspaper reader the same help that he could get from a monocle.

The invention of lenses was a necessary preliminary to the invention of the telescope, for, as Huygens remarked, it would require a superhuman genius to make the invention theoretically.

The retina of the eye on which the image is to be received has structure. We may compare the picture on the retina to a design embroidered in wool-work, which also has a structure. Clearly such a design cannot embody details which are smaller than the mesh of the canvas which is to carry the coloured stitches. The only way to get in more details is to make the design, or rather such diminished part of it as the canvas can accommodate, on a larger scale. Similarly with the picture on the retina. The individual rods and cones correspond with the individual meshes of the canvas. If we want more detail of an object we must make the picture on the retina larger, with the necessary sacrifice of the field of view. If the object is distant, we want for this a lens of longer focus instead of the eye lens. We cannot take the eye lens away, but, what amounts to nearly the same thing, we can neutralise it by a concave lens of equal power put right up to it, called the eye-piece. Then we are free to use a long focus lens called the telescopic objective to make a larger picture on the retina. It must of course be put at the proper distance out to make a distinct picture. This is a special case of the Galilean telescope which lends itself to simple description. It is of no use to make the picture larger if we lose definition in the process. The enlarged image must remain sharp enough to take advantage of the fine structure of the retinal screen that is to receive it. It will not be sharp enough unless we make the lens of greater diameter than the eye. Another reason for using a large lens is to avoid a loss of brightness.

It seems paradoxical that the image of a star should be smaller the larger the telescope. Nevertheless it is a necessary result of the wave character of light. We cannot see the true nature of, for example, a double star, unless the two images are small enough not to overlap and far enough apart to fall on separated elements of the observer's retina.

When the problem is to examine small objects we look at them as close as we can: here the short-sighted observer has an advantage. By adding a lens in front of the

eye lens to increase its power we can produce a kind of artificial short sight and get closer than we could otherwise, so that the picture on the retina is bigger. This is a simple microscope and we can use it to examine the image produced by an objective lens; if this image is larger than the object under examination we call the whole arrangement a compound microscope.

Given perfect construction there is no limit in theory to what a telescope can do in revealing distant worlds. It is only a question of making it large enough. On the other hand, there is a very definite limit to what the microscope used with, say, ordinary daylight can do. It is not that there is any difficulty in making it magnify as much as we like. This can be done, *e.g.*, by making the tube of the microscope longer. The trouble is that beyond a certain point magnification does no good. Many people find this a hard saying, but it must be remembered that a large image is not necessarily a good image. We are up against the same difficulty as before. A point on the object is necessarily spread out into a disc in the image, due to the coarseness of structure of light itself as indicated by its wave-length. I cannot go into the details, but many of you will know that points on the object which are something less than half a wave-length or say a one hundred-thousandth of an inch apart, cannot be distinctly separated. This is the theoretical limit for a microscope using ordinary light, and it has been practically reached. The early microscopists would have thought this more than satisfactory; but the limit puts a serious obstacle in the way of biological and medical progress to-day. For example, the pathogenic bacteria in many cases are about this size or less; and there is special interest in considering in what directions we may hope to go further.

Since microscopic resolution depends on having a fine structure in the light itself, something though not perhaps very much, may be gained by the use of ultra-violet light instead of visible light. It then becomes necessary to work by photography. We are nearing the region of the spectrum where almost everything is opaque. In the visual region nearly every organic structure is transparent and to get contrast stains have to be used which colour one part more deeply than the other. In the ultra-violet, on the other hand, we get contrast without staining and, as Mr. J. W. Barnard has shown, the

advantage lies as much in this as in the increased resolving power. For example, using the strong ultra-violet line of the mercury vapour lamp, which has about half the wave-length of green light, he finds that a virus contained within a cell shows up as a highly absorptive body in contrast with the less absorptive element of the cell. So that ultra-violet microscopy offers some hope of progress in connection with this fundamental problem of the nature of viruses.

With ultra-violet microscopy we have gone as far as we can in using short waves with ordinary lenses made of matter, for the available kinds of matter are useless for shorter waves than these, and it might well seem that we have here come to a definite and final end. Yet it is not so. There are two alternatives, which we must consider separately. Pradoxical as it may seem, for certain radiations we can make converging lenses out of empty space; or alternatively we can make optical observations without any lenses at all.

The longstanding controversy which raged in the nineties of the last century as to whether cathode rays consisted of waves or of electrified particles was thought to have been settled in favour of the latter alternative. But scientific controversies, however acutely they may rage for a time, are apt, like industrial disputes, to end in compromise; and it has been so in this instance. According to our present views the cathode rays in one aspect consist of a stream of electrified particles; in another, they consist of wave trains, the length being variable in inverse relation to the momentum of the particles.

Now cathode rays have the property of being bent by electric or magnetic forces, and far-reaching analogies have been traced between this bending and the refraction of light by solids; indeed, a system of 'electron optics' has been elaborated which shows how a beam of cathode rays issuing from a point can be reassembled into an image by passing through a localised electrostatic or magnetic field having axial symmetry. This constitutes what has been called an electrostatic or magnetic lens. It is then possible to form a magnified image of the source of electrons on a fluorescent screen, and that is the simplest application. But we can go further and form an image of an obstructing object such as a fine wire by means of one magnetic lens, acting as objective, and amplify it by means of a second magnetic lens, which is

spoken of as the eye-piece, though of course it is only such by analogy, for the eye cannot deal directly with cathode rays. The eye-piece projects the image on to a fluorescent screen, or photographic plate. So far we have been thinking of the electron stream in its corpuscular aspect. But we must turn to the wave aspect when it comes to consideration of theoretical resolving power. The wave-length associated with an electron stream of moderate velocity is so small that if the electron microscope could be brought to the perfection of the optical microscope, it should be able to resolve the actual atomic structure of crystals. This is very far indeed from being attained, the present electron microscope being much further from its own ideal than were the earliest optical microscopes. Nevertheless experimental instruments have been constructed which have a resolving power several times better than the modern optical microscope. The difficulty is to apply them to practical biological problems.

It is not to be supposed that the histological technique so skilfully elaborated for ordinary microscopy can at once be transferred to the electron microscope. For example, the relatively thick glass supports and covers ordinarily used are out of the question. Staining with aniline dyes is probably of little use, and the fierce bombardment to which the delicate specimen is necessarily exposed will be no small obstacle. Certain standard methods, however, such as impregnation with osmium, seem to be applicable: and there is some possibility that eventually the obscure region between the smallest organisms and the largest crystalline structure may be explored by electron microscopy.

In referring to the limitations on the use of lenses I mentioned the other alternative that we might, in order to work with the shortest waves, dispense with lenses altogether: and in fact in using X-rays this is done. We are then limited to controlling the course of the rays by means of tubes or pinholes. This restriction is so serious that it altogether defeats the possibility of constructing a useful X-ray microscope analogous to the optical or the electron microscope. In spite of this the use of X-rays is of fundamental value for dealing with a particular class of objects, namely, crystals which themselves have a regular spacing, comparable in size with the length of the waves. Just as the spacing of

a ruled grating (say one $1/20,000$ th of an inch) can be compared with the wave-length of light by measuring the angle of diffraction, so the spacing of atoms in a crystal can be compared with the wave-length of X-rays. But here the indications are less direct than with the microscope, and depend on the object having a periodic structure. So that the method hardly falls within the scope of this address. How essential the difference is will appear if we consider that the angle to be observed becomes greater and not less the closer the spacing of the object under test.

Colour vision is one of nature's most wonderful achievements, though custom often prevents our perceiving the wonder of it. We take it for granted that anyone should readily distinguish the berries on a holly bush, and we are inclined to be derisive of a colour-blind person who cannot do so. But so far anatomy has told us little or nothing of how the marvel is achieved. Experiments on colour vision show that three separate and fundamental colour sensations exist. It is probable that the cones of the retina are responsible for colour vision and the rods for dark adapted vision which does not discriminate colour. But no division of the cones into three separate kinds corresponding to the three colour sensations has ever been observed. Nor is any anatomical peculiarity known which allows a colour-blind eye to be distinguished from a normal one.

Can artificial resources help to improve colour discrimination? In some interesting cases they can. Indeed, the whole subject of spectroscopy may be thought of as coming under this head. We can recognise the colour imparted by sodium to a flame without artificial help. When potassium is present as well, the red colour due to it can only be seen when we use a prism to separate the red image of the flame from the yellow one. Such a method has its limitations, because if the coloured images are more numerous they overlap, and the desired separation is lost. To avoid this it is necessary to make a sacrifice, and to limit the effective breadth of the flame by a more or less narrow slit. And if the images are very numerous the slit has to be so narrow that all indication of the breadth of the source is lost. This, of course, is substantially the method of spectroscopy, into which I do not enter further. But there is an interesting class of cases where we cannot afford to sacrifice the

form of the object entirely to colour discrimination. Consider, for example, the prominences of the sun's limb, which are so well seen against the darkened sky of an eclipse, but are altogether lost in the glare of the sky at other times. In order to see them prismatic dispersion is made use of, and separates the monochromatic red light of hydrogen from the sky background. A slit must be used to cut off the latter: but if it is too narrow the outlines of the prominence cannot be seen. By using a compromise width it is possible to reconcile the competing requirements in this comparatively easy case. Indeed, M. B. Lyot, working in the clear air of the observatory of the Pic du Midi, where there is less false light to deal with, has even been able to observe the prominences through a suitable red filter, which enables the whole circumference of the sun to be examined at once, without the limitations introduced by a slit. A much more difficult problem is to look for bright hydrogen eruptions projected on the sun's disc, and at first sight this might well seem hopeless. A complete view of them was first obtained by photography, but I shall limit myself to some notice of the visual instrument perfected by Hale and called by him the spectrohelioscope. A very narrow slit has to be used, and hence only a very small breadth of the sun's surface can be seen at any one instant. But the difficulty is turned by very rapidly exposing to view successive strips of the sun's surface side by side. The images then blend, owing to persistence of vision, and a reasonably broad region is included in what is practically a single view. I must pass over the details of mechanism by which this is carried out.

There are now a number of spectrohelioscopes over different parts of the world, and a continuous watch is kept for bright eruptions of the red hydrogen lines. Already these are found to be simultaneous with the 'fading' of short radio waves over the illuminated hemisphere of the earth, and the brightest eruptions are simultaneous with disturbances of terrestrial magnetism. At the Mount Wilson Observatory such eruptions have been seen at the same time at widely separated points on the sun, indicating a deep-seated cause. There are therefore very interesting and fundamental questions within the realm of this method of investigation.

We have so far been mainly considering how

we may adapt our vision for objects too small or too far off for unassisted sight, and for colour differences not ordinarily perceptible. This is chiefly done by supplementing the lens system of the eye by additional lenses or by prisms. We cannot supplement the retina, but in certain cases we can do better. We can substitute an artificial sensitive surface which may be either photographic or photoelectric.

That certain pigments are bleached by light is an observation that must have obtruded itself from very early times—indeed, it is one of the chief practical problems of dyeing to select pigments which do not fade rapidly. If a part of the coloured surface is protected by an opaque object—say a picture or a mirror hanging over a coloured wallpaper—we get a silhouette of the protecting object, which is in essence a photograph.

Again, it is a matter of common observation that the human skin is darkened by the prolonged action of the sun's light, and here similarly we may get what is really a silhouette photograph of a locket, or the like, which protects the skin locally. In this case we are perhaps retracing the paths which Nature herself has taken: for the evolution of the eye is regarded as having begun with the general sensitiveness to light of the whole surface of the organism.

The sensitivity of at all events the dark adapted eye depends on the accumulation on the retinal rods of the pigment called the visual purple of which the most striking characteristic is its ready bleaching by light. We can even partially 'fix' the picture produced in this way on the retina of, for example a frog by means of alum solution. This brings home to us how clearly akin are the processes in the retina to those in the photographic plate, even though the complexity of the former has hitherto largely baffled investigation.

There are then many indications in nature of substances sensitive to light, and quite a considerable variety of them have from time to time been used in practical photographic processes. But compounds of silver, which formed the basis of the earliest processes, have maintained the lead over all others. The history of photography by means of silver salts cannot be considered a good example of the triumph of the rational over the empirical. For instance, the discovery of developers came about thus.

The first workers, Wedgewood and Davy (1802), had found that they got greater sensitivity by spreading the silver salt on white leather instead of paper. An early experimenter, the Rev. J. B. Reade (1837), was anxious to repeat this experiment, and sacrificed a pair of white kid gloves belonging to his wife for the purpose. When he wished to sacrifice a second pair, the lady raised a not unnatural objection, and he said, 'Then I will tan paper.' He treated paper with an infusion of oak galls and found that this increased the sensitivity greatly. It amounted to what we should call exposing and developing simultaneously. But, in using the method, it is easily observed that darkening continues after exposure is over, and this leads to beginning development after the exposure. This step was taken by Fox Talbot a year or two afterwards. Instead of crude infusion of galls he used gallic acid. Later pyrogallie acid was used instead of gallic acid, and still survives.

The use of gelatine as a medium to contain the silver halide was a more obvious idea. But it was not so easy to foresee that the sensitivity of silver salts would be much further increased when they were held in this medium. For long this remained unexplained, until it was noticed that some specimens of gelatine were much more active than others. This was ultimately traced by S. E. Sheppard to the presence of traces of mustard oil, a sulphur compound, in the more active specimens. This, in turn, depends in all probability on the pasturage on which the animals that afford the gelatine have been fed. The quantity present is incredibly small, comparable in quantity with the radium in pitchblende.

The value to science as well as to daily life of the gelatine dry plate or film can hardly be over-estimated. Take, for instance, the generalised principle of relativity, which attempts with considerable success to reduce the main feature of the cosmical process to a geometrical theory. The crucial test requires us to investigate the gravitational bending of light, by photographing the field of stars near the eclipsed sun. For this purpose the gelatine dry plate has been essential: and here, as we have seen, we get into complicated questions of bio-chemistry. This is to my mind a beautiful example of the interdependence of different branches of science and of the disadvantages of undue specialisation (or should I say generalisa-

tion?). We may attempt to reduce the cosmos to the dry bones of a geometrical theory, but in testing the theory we are compelled to have recourse again to the gelatine which we have discarded from the dry bones!

To come back, however, to the development of the photographic retina, as I may call it. As is well known, the eye has maximum sensitivity to the yellow-green of the spectrum, but ordinary silver salts are not sensitive in this region. Their maximum is in the blue or violet, and ranges on through ultra-violet to the X-ray region. It was not at all easy to extend it on the other side through green, yellow and red to infra-red. The story of how this was ultimately attained is one more example in the chapter of accidental clues skilfully followed up which forms the history of this subject.

In 1873, Dr. Hermann Vogel, of Berlin, noticed that certain collodion plates of English manufacture, which he was using for spectrum photography, recorded the green of the spectrum to which the simple silver salts are practically insensitive. The plates had been coated with a mixture which contained nitrate of uranium, gum, gallic acid and a yellow colouring matter. What the purpose of this coating was is not very obvious. It rather reminds one of mediæval medical prescriptions which made up in complexity what they lacked in clear thinking. But Vogel concluded with true scientific insight that it must owe the special property he had discovered to some constituent which absorbs the green of the spectrum more than the blue: for conservation of energy required that the green should be absorbed if it is to act on the plate. He then tried staining the plate with coralline red, which has an absorption band in the green, with the expected result. With much prescience he says: 'I think I am pretty well justified in inferring that we are in a position to *render bromide of silver sensitive for any colour we choose*. Perhaps we may even arrive at this, namely photographing the ultra-red as we have already photographed the ultra-violet.' It was, however, half a century before this far-seeing prophecy was fully realised. The development of the aniline colour industry gave full scope for experiment, but it has been found by bitter experience that dyes which can produce the colour sensitiveness are often fatal to the clean working and keeping qualities of the

plate. However, success has been attained, largely by the efforts of Dr. W. H. Mills, of the Chemical Department of this University, and of Dr. Mees, of the Kodak Company; and we all see the fruits of it in the photographs by lamplight which are often reproduced in the newspapers.

It is now known in what direction the molecular structure of the sensitising dye must be elaborated in order to push the action further and further into the infra-red, and the point when water becomes opaque has nearly been reached, with great extension of our knowledge of the solar spectrum. The spectra of the major planets have also been extended into the infra-red and this has given the clue as to the true origin of the mysterious absorption bands due to their atmospheres which had baffled spectroscopists for more than a generation. These bands have been shown by Wildt to be due to methane or marsh gas. Neptune, for example, has an atmosphere of methane equivalent to 25 miles thickness of the gas under standard conditions. In this Neptunian methane we have a paraffin certainly not of animal or vegetable origin; and I venture in passing to make the suggestion that geologists might usefully take it into consideration in discussing the origin of terrestrial petroleum.

The photographic plate is not the only useful substitute for the human retina. We have another in the photoelectric surface. The history of this discovery is of considerable interest. Heinrich Hertz, in his pioneering investigation of electric waves (1887), made use of the tiny spark which he obtained from his receiving circuit as an indicator. The younger part of my audience must remember that this was before the days of valves and loud speakers. His experiments were done within the walls of one room. When he boxed in the indicating spark so as to shield it from daylight and make it easier to see, he found that this precaution had exactly the opposite effect—the spark became less instead of more conspicuous. To express it shortly and colloquially, this action was found to depend on whether or not the spark of the receiver could see the spark of the oscillator. Moreover, seeing through a glass window would not do. It was ultra-violet light from the active spark that influenced the passive spark. Further, Hertz was able to determine that the action occurred

mainly, if not entirely, at the cathode of the passive spark.

The next step was taken by Hallwachs, who showed that it was not necessary to work with the complicated conditions of the spark. He found that a clean zinc plate negatively charged rapidly lost its charge when illuminated by ultra-violet light.

The final important step was in the use of a clean surface of alkali metal *in vacuo* which responds to visible light and passes comparatively large currents. This constitutes the photoelectric cell very much as we now have it, and was due to two German school-masters J. Elster and H. Geitel. English physicists who met them during their visit to Cambridge a generation ago will not fail to have agreeable memories of their single-minded enthusiasm and devoted mutual regard. Sir J. J. Thomson has recalled them to our recollection in his recent book. They could scarcely have foreseen that their work, carried out in a purely academic spirit, would make possible the talking films which give pleasure to untold millions.

The sensitiveness of the dark-adapted eye has often been referred to as one of its most wonderful features; but under favourable conditions, the sensitivity of a photoelectric surface may even be superior. According to our present ideas, no device conceivable could do more than detect every quantum which fell upon it. Neither the eye nor the photoelectric surface comes very near to this standard, but it would seem that the falling short is rather in detail than in principle. The action of the photoelectric cell depends on the liberation of an electron by one quantum of incident energy, and under favourable conditions the liberation of one electron can be detected, by an application of the principle of Geiger's counter. The action of the dark-adapted eye depends on the bleaching of the visual purple. According to the results of Dartnall, Goodeve and Lythgoe it appears likely that one quantum can bleach a molecule of this substance, and in all probability this results in the excitation of a nerve fibre, which carries its message to the brain.

The photoelectric cell can be used like the photographic plate at the focus of an astronomical telescope. It might seem from the standpoint of evolution a retrograde step to substitute a single sensitive element for the 137 million such elements in the human eye. In this connection it is interesting to note

that in certain invertebrate animals eyes are known which have the character of a single sensitive element, with a lens to concentrate the light upon it. Such an eye can do little more than distinguish light from darkness. But its artificial counterpart using the photoelectric surface has the valuable property that the electric current which indicates that light is falling upon it can be precisely measured, so as to determine the intensity of the light. In contrast with photographic action, the energy available to produce the record comes not from the original source of light, which only, as it were, pulls the trigger, but from the battery in the local circuit, and it may be amplified so as to actuate robust mechanisms. It has been applied with success to guiding a large telescope, or in a humbler sphere, to open doors, or even to catch thieves.

However, the scientific interest lies more in the possibility of accurate measurement. As an interesting example we might take the problem of measuring the apparent diameter of the great nebula in Andromeda. As is known, modern research tends to indicate that the Andromeda nebula and other like systems are the counterparts of the galaxy, being in fact island universes. But until lately there was a serious difficulty in that all such systems appeared to be considerably smaller than the galaxy. Stebbins and Whitford, by traversing a telescope armed with a photoelectric cell across the nebula, have found that its linear dimensions were twice as great as had been supposed, reducing the discrepancy of size to comparatively little.

But, it may be suggested, could we not go further and make a photoelectric equivalent, not only for the rudimentary kind of eye which has only a single sensitive element, but for the developed mammalian eye which has an enormous number? Could we not build up on separated photoelectric elements a complete and detailed picture? In point of fact this has been done in the development of television; and since this new art which interests us all can properly be considered as an extension of the powers of normal vision, no excuse is needed for devoting some consideration to it. We must divide the photoelectric surface into minute patches which are electrically insulated from one another. This is not too difficult; but if it were proposed directly to imitate nature, and attach a wire, representing a nerve fibre,

to each of these patches, so as to connect it to the auxiliary apparatus, we might well despair of the task; for there are probably half a million such connections between the human retina and the brain. In the artificial apparatus for television, one single connection is made to serve, but it is in effect attached to to each of the patches in rapid succession by the process of 'scanning' the image. The photoelectric mosaic is on one side of a thin mica sheet, and a continuous metal coating on the other side gives the connection, which is by electrostatic induction. Each element of the surface forms a separate tiny condenser with the opposing part of the back plate. Scanning is achieved by rapidly traversing a beam of electrons over the mosaic line by line. The whole surface, and therefore each element, must be scanned at least twenty times a second. In the intervals an element is losing electrons more or less rapidly. The scanning beam comes along, and restores the lost electrons, discharges the little condenser found by the element and the back plate and sends an electric signal into the wire attached to this plate. The strength of this signal will depend on how many electrons the element had lost since the previous scanning, and thus on the luminous intensity of that part of the image. An important point is that the element is in action all the time, and not only while it is individually being scanned. We have thus transmuted the momentary picture into a series of electric pulses occupying in all a time of one-twentieth of a second and these can be amplified and sent out as wireless signals. How are they to be turned back again into a visible picture at the other end? Well, that is not perhaps so difficult as the first conversion of the picture into signals. We must make a beam of electrons follow and imitate the periodic movements of the scanning beam at the other end. The beam of electrons falls on a luminescent screen, and makes it light up, more or less brightly according to the intensity of the electron beam. If we use the incoming signals to modulate the electron beam, we can make them correspond with the intensities at the sending end, and the original picture is reconstructed piece by piece. The reconstruction is completed in one-twentieth of a second or less, and the process begins again. The successive pictures blend into one another as in the cinema, and movement is shown with apparent continuity.

It seems not unlikely that the electric

eye or iconoscope, as it has been called, may have applications apart from television. Dr. V. K. Zworykin, who took an important part in its development, suggested that it might be used to make visible the image in the ultra-violet microscope, which would be much too faint for direct projection on a fluorescent screen. For that purpose the sending and receiving apparatus would, of course, be connected directly, without radio transmission. It might also be used for rapid photography, if the photographic plate replaced the viewing screen. The beauty of the device is that the energy is supplied locally, the distant light source merely releasing it. The principle of amplification may thus perhaps be applied to the photographing of faint objects.

I come to the close of this part of my subject.

Much of modern scientific doctrine appears at first sight to have an elusive and even metaphysical character, and this aspect of it seems to make the strongest appeal to many cultivated minds. Yet upon the whole, the main triumphs of science lie in the tangible facts which it has revealed; and it is these which will without doubt endure as a permanent memorial to our epoch. My main thesis has been that these are discovered by methods not essentially different from direct scrutiny. It is hoped that the present survey may remind you that if we allow for a reasonable broadening of the original meaning of the words, it remains true after all that 'seeing is believing'.

II.

SCIENCE AND WARFARE.

During the Great War itself, few scientific men in any country doubted that it was their duty to do what they could do to apply their specialised knowledge to the purposes of war; nor was it often suggested by publicists that there was any countervailing consideration: on the contrary they urged strongly that our resources in this direction should be efficiently mobilised. It is chiefly in vague general discussions that the opposite view becomes vocal.

Science, it is urged, is the source of all the trouble: and we may look to scientific men for some constructive contribution to finding a remedy. It is worth while to inquire what basis there is for this indictment, and whether, in fact, it is feasible for

men of science to desist from labours which may have a disastrous outcome, or at any rate to help in guiding other men to use and not to abuse the fruits of those labours. I may say at the outset that I have no sanguine contribution to make. I believe that the whole idea that scientific men are specially responsible is a delusion born of imperfect knowledge of the real course of the process of discovery. Indeed, very much the same complaint was made before the scientific era. Let me refer you to Shakespeare's play of *Henry IV*:—

' Great pity, so it was
This villainous saltpetre should be digged
Out of the bowels of the harmless earth
Which many a good tall fellow had destroyed
So cowardly.'

The quotation leads us to inquire how far the further development of this particular kind of frightfulness into modern high explosives was deliberate or not.

In the course of systematic study of the chemistry of carbon compounds it was inevitable that the action of nitric acid on substances like benzene, toluene, glycerine, cellulose and the like should be tried. No one could foresee the result. In the case of benzene, we have nitrobenzene, the key to the aniline dye industry. In the case of glycerine, Sobrero obtained in 1846 the highly explosive liquid called nitro-glycerine. He meant no harm, and in fact his discovery lay dormant for many years, until Nobel turned his attention to the matter in 1863, and showed how by mixing nitro-glycerine with other substances, solid explosives could be made which admitted of safe handling. Dynamite was one of them. They proved invaluable in the arts of peace, *e.g.*, in mining and in making railway tunnels, such as those through the Alps. They were used by the Irish Fenians in the dynamite outrages of the eighties. These attempted outrages were not very successful, and so far as I know no one was inclined to blame science for them, any more than for the Gunpowder Plot. Like the latter, they came to be considered slightly comic. If any one doubts this, he may agreeably resolve his doubts by reading R. L. Stevenson's story *The Dynamiter*. At all events, high explosives had been too long in use in peaceful industry for their misuse to be laid directly to the account of science.

Coming next to poison gas. We read that Pliny was overwhelmed and killed by sulphur dioxide in the eruption of Vesuvius in A.D. 79. During the Crimean War, the veteran admiral Lord Dundonald urged that the fumes of burning sulphur should be deliberately used in this way, but the suggestion was not adopted. Even if it had been, scientific research *ad hoc* would obviously have had little to do with the matter. During the Great War, chlorine was used on a large scale. I need hardly insist that chlorine was not isolated by chemists for this purpose. It was discovered 140 years before, as a step in the inquiry into the nature of common salt.

Coming to the more recondite substances, we may take mustard gas—really a liquid—as typical. It is much more plausible to suggest that here was a scientific devilment, deliberately contrived to cripple and destroy. But what are the real facts?

Referring to Watt's *Dictionary of Chemistry* (edition of 1894), there is an article of less than forty words about mustard gas (under the heading of dichloro-diethyl sulphide). After the method of preparation used by Victor Meyer has been mentioned, the substance is dismissed with the words 'oil, very poisonous and violently inflames the skin. Difference from diethyl sulphide.'

There are sixteen other compounds described at comparable length on the same page. So far as I know, none of them is of any importance. A not uncommon type of critic would probably say that the investigation of them had been useless, the work of impractical dreamers, who might have been better employed. One of these substances, namely mustard gas, is quite unexpectedly applied to war, and the production of it is held by the critics to be the work not of dreamers, but of fiends whose activities ought to be suppressed! Finally at the bottom of the page begins a long article on chloroform. This substance, as you know, has relieved a great deal of pain, and on the same principle the investigator who produced it was no doubt an angel of mercy. The trouble is that all the investigators proceeded in exactly the same spirit, the spirit that is of scientific curiosity, and with no possibility of telling whether the issue of their work would prove them to be fiends, or dreamers, or angels.

Again, there is the terror of thermite incendiary bombs, spreading fire broadcast through our great cities. The notion is sometimes encountered that thermite was invented for this purpose. Nothing could be further from the truth. I first made acquaintance with it myself in 1901 by hearing a lecture at the Royal Institution by the late Sir William Roberts Austen on 'Metals as Fuel'.¹ He drew attention to the great amount of energy which was liberated when aluminium combined with oxygen, and showed how aluminium powder mixed with red oxide of iron would react violently with it, withdrawing the oxygen from the iron, and becoming brilliantly incandescent in the process. He showed further how this mixture, called thermite, could be used for heating metal work locally, so as to make welds, *e.g.*, in joining two iron pipes end to end. I venture to say that it never occurred to him or to any of his hearers that thermite had any application in war.

In discussions of this kind a distinction is often implied between what I may call old-fashioned knowledge and modern scientific knowledge. The latter is considered to be the special handmaid of 'frightfulness'. The futility of this distinction is easily seen by considering a special case. Iron is thought of as belonging to the pre-scientific era, while aluminium is thought to belong to the scientific era. From the standpoint of chemistry both are metals, and the problem of producing them in either case is a chemical one. When produced they both have their function in 'frightfulness'; iron to cut and stab; aluminium to make thermite bombs to burn and destroy. If modern science makes its contribution to 'frightfulness' in giving us aluminium, ancient craft did so in giving us iron. It is obviously absurd to make any distinction in principle between the two cases. Science properly understood includes all real knowledge about material things, whether that knowledge is old or new.

All these terrors have only become applicable against a civilian population by the development of aircraft. Military objects

were certainly not the incentive of the successful pioneers of artificial flight. They were fascinated at first by the sport of gliding, and afterwards by a mechanical transport problem.

It is true that brilliant writers of imaginative fiction, such as Jules Verne and H. G. Wells, had foretold all, and more than all, the horrors that have since come to pass. But it is perhaps more to the point to inquire what were the contemporary views of practical men. The Wrights made their first successful flight in 1903. In 1904 I myself heard the then First Sea Lord of the Admiralty repudiate with scorn the suggestion that the Government were interesting themselves in the matter; and I know with equal definiteness that even as late as 1908 the Chief of the Imperial General Staff did not believe in the military importance of flight. Would it be fair then to blame the inventors for not having realised it, and for not having stayed their hands?

Summing up what may be learnt from the experience of the past, I think we may say that the application of fundamental discoveries in science to purposes of war is altogether too remote for it to be possible to control such discoveries at the source.

For good or ill, the urge to explore the unknown is deep in the nature of some of us, and it will not be deterred by possible contingent results, which may not be, and generally are not, fully apparent till long after the death of the explorer. The world is ready to accept the gifts of science, and to use them for its own purposes. It is difficult to see any sign that it is ready to accept the advice of scientific men as to what those uses should be.

Can we then do nothing? Frankly, I doubt whether we can do much, but there is one thing that may be attempted. The Association has under consideration a division for study of the social relations of science which will attempt to bring the steady light of scientific truth to bear on vexed questions. We rejoice to know that our distinguished American visitors are in sympathy with this aim, and we hope that our discussions with them will bear useful, if modest, fruit in promoting international amity.

¹ *Proc. R.I.*, 1901, Feb. 23, 16, 496.