

radioactivity ($2.5 \pm 0.05 \beta/c/g/min$) than the khondalites ($0.5 \pm 0.05 \beta/c/g/min$) due to the presence of monazite as an accessory mineral⁷. Thorium strongly concentrates in acidic igneous rocks¹⁵ and its content in them is around 13 g/ton¹⁶. The concentration of thorium continues during pegmatitic stage even after the close of the main stage of crystallization¹⁵. Hence it is concluded that the monazites are derived from charnockites and pegmatites which occur in the drainage basin of the hinterland.

The concentration of monazite in considerable proportions and its high thorium content calls for an evaluation of the commercial potential for exploitation along with other associated minerals like ilmenite, garnet, sillimanite and zircon from this area, keeping in view among other uses, also the requirement of thorium in fast breeder reactors.

1. Tipper, G. H., *Rec. Geol. Surv. India*, 1994, **44**, 195.
2. Mahadevan, C. and Sriramadas., A., *Proc. Indian Acad. Sci.*, 1948, **27**, 275-278.
3. Ramamohana Rao, T., Shanmukha Rao, Ch. and Sanyasi Rao, K., *J. Geol. Soc. India*, 1982, **23**, 284-289.
4. Reddy, D. R. S., Varma, D. D., Reddy, K. S. N. and Prasad, V. S. S., in abstract volume of XII Convention of Indian Association of Sedimentologists, Goa University, 20-22 Dec. 1995, p. 53.
5. Rajasekhar Reddy, D. and Siva Sankara Prasad, V., Paper communicated to *Indian Minerals*.

6. Dhanunjaya Rao, G., Krishnaiah Setty, B. and Raminaidu, Ch., in *Exploration and Research for Atomic Minerals*, 1989, vol. 2, pp. 147-155.
7. Rao, A. T. and Prakasa Rao, Ch. S., *Indian J. Mar. Sci.*, 1980, **9**, 214-216.
8. Deer, W. A., Howie, R. A. and Zussman, J., in *Rock Forming Minerals*, Longman, London, 1975, vol. 5, pp. 339-346.
9. Bowie, S. H. U. and Horne, J. E. T., *Min. Mag.*, 1953, **30**, 93.
10. Kamineni, D. C., Rao, A. T. and Bonardi, M., *Mineral Petrol.*, 1991, **45**, 119-130.
11. Murata, K. J., Dutra, C. V., Da Coasta, M. T. and Branco, J. J. R., *Geochim. Cosmochim. Acta*, 1959, **16**, 1-14.
12. Kshira Sagar, T. V. S. R. and Nagamalleswara Rao, B., *Indian Mining Eng. J.*, 1993, **XLI**, 31-33.
13. Murty, M. S., *Curr. Sci.*, 1958, **27**, 347-348.
14. Mahadevan, C. and Sathapathi, *Curr. Sci.*, 1948, **17**, 297.
15. Rankama, K. and Sahama, Th. G., in *Geochemistry*, John Wiley, New York, 1950, p. 571.
16. Evans, R. D. and Goodman, C., *Bull. Geol. Soc. Am.*, 1941, **52**, 459.

ACKNOWLEDGEMENTS. The present work is a part of a DST, New Delhi-sponsored project. We are thankful to Mr. Robert Mac Kay, Scanning Electron Microprobe Image Analysis Laboratory, Department of Earth Sciences, Dalhousie University, Canada for carrying out the microprobe analyses. Our sincere thanks are due to the anonymous referee for suggestions.

Received 28 July 1997; revised accepted 15 October 1997

Erratum

Observed and theoretical acceleration response spectra in the Tehri region: Implications for the seismic hazard in the region

K. N. Khattri

(*Curr. Sci.*, 1995, **69**, 161-171)

Due to an inadvertent oversight, Table 1 showing the velocity and Q model is in error. The correct model is given below.

Table 1. Velocity and Q model

Thickness (km)	V _p (km/s)	Q _p	V _s (km/s)	Q _s	Density (g/cc)
0.05	0.9	10	0.5	5	1.8
0.5	1.75	30	1.0	15	1.8
2.0	4.0	100	2.8	50	2.4
14.0	5.2	4000	2.97	2000	2.6
30.0	6.0	4000	3.43	2000	2.9
Infinity	8.33	1000	4.83	500	3.3