

How do we 'miss the wood for the trees'? Do females have natural flair for lullaby? Cognitive neuroscience tells it all (or does it?)

T. Ramakrishna

Roger Sperry was awarded the 1981 Nobel Prize in Physiology or Medicine for his work on split-brain patients. He showed that in patients whose left and right cerebral hemispheres have been separated (by cutting the 200 million fibres which make up the corpus callosum that connects both the hemispheres), it is the left cerebral hemisphere that processes language whereas the right cerebral hemisphere processes the objects. It was possible to test the functions of each hemisphere




Figure 1. S and V represent global level and F and P, the local level.




Figure 2. The cup made up of small anchors and the anchor made up of small cups constitute the global level. Small cups within the anchor and the small anchors within the cup constitute the local level.

separately in these patients by showing pictures of objects either through left visual field, thereby feeding this information only to the right cerebral hemisphere (because of the crossing over of the visual information pathways to the contra lateral hemisphere) or vice versa. So long as the picture of any object is registered in the left cerebral hemisphere, the patient would be able to tell the name of the object. However if the picture of the same object is registered in the right cerebral hemisphere, the patient would be unable to tell the name of the object. Through several such ingenious experiments, Sperry was successful in proving that in the majority of the right handers language processing is carried out in the left hemisphere whereas the object processing is done in the right cerebral hemisphere.

This epoch-making discovery in cognitive neuroscience has come to be known as lateralization of hemispheric mechanisms/functions¹.

Cognitive neuroscience has come a long way since the celebrated discovery of Sperry as is evidenced by a special issue of *Science* on cognitive neuroscience very recently². While in the past most neuroscientists focused on individual brain regions and their responses, it is now becoming clear that the processing of information that leads to complex behaviours such as learning and memory involves multiple brain regions operating in interactive and integrated synchrony. The methodology such as the measurement of relative regional cerebral blood flow (rCBF) by recording the regional distribution of cerebral radioactivity following the intravenous injection of ¹⁵O-labelled water, known as positron emission tomography (PET scanning), and the functional magnetic resonance imaging (fMRI), made it possible for researchers to examine the responses of distributed brain networks while an individual is performing a task.

Armed with these powerful tools, neuroscientists have given us new knowledge bearing on the questions: (i) How does the brain process global information versus local information, with reference to letters (language) and objects (pictures), and (ii) whether there is any gender-based difference in the brain mechanism(s) for language processing. The first question was dealt with by Fink and his associates at the Institute of Neurology, London and at the Radcliff Infirmary, Oxford^{3,4}, while the second question was addressed by Shaywitz and his colleagues at the University School of Medicine, New Haven, Connecticut⁵.

The perceptual world is organized hierarchically (the forest consists of trees and the trees consist of leaves and so on) and the visual attention can emphasize either the overall picture (global form) or the details within (local components). For example, the letters S and V (Figure 1) used in the experiments by Fink and his colleagues represent the global level and F and P, the local level. Their findings indicate that globally-directed attention involves the right hemisphere and locally-directed attention, the left hemisphere. More specifically, they demonstrated lateralized neural activity in the left prestriate cortex during local processing and in the right prestriate cortex during global processing. In these experiments they used letter-based hierarchical stimuli³. In another experiment, they used object-based hierarchical stimuli (Figure 2), wherein, the cup made up of small anchors and the anchor made up of small cups constitute the global level. Small cups within the anchor and the small anchors within the cup constitute the local level. There is now greater right-sided activation of prestriate cortex for local processing and greater left-sided activation for global processing which is the opposite of that seen with the letter-based stimuli of the earlier experiment! The stimulus category (whether letter or

INFORMATION FOR CONTRIBUTORS

GENERAL

All manuscripts should be addressed to the Editor, *Current Science*, P. B. No. 8001, C. V. Raman Avenue, Bangalore 560 080. Submission of an article will be held to imply that it has not been previously published and is not under consideration for publication elsewhere; and further, that if accepted, it will not be published elsewhere. *Three copies of contributions of all categories* are required, with a letter of transmittal giving (i) names and complete addresses of the authors and (ii) title of the contribution and the category in which it is submitted (see below).

Current Science is a multidisciplinary journal and therefore research and review papers of general significance that are written clearly and well organized will be given preference. All papers will be first assessed by a Reviewing Editor. Papers found unsuitable in terms of the overall requirements of the journal will be returned to the authors. The others will be sent for detailed review. *Both solicited and unsolicited material will be reviewed.* Authors of these papers will be notified of acceptance, rejection, or need for revision of the paper. Returned papers cannot be resubmitted.

Illustrations and other material to be reproduced from other publications must be properly credited; it is the authors' responsibility to obtain permission for reproduction (copies of letters of permission should be sent).

CATEGORIES OF MANUSCRIPT

General articles (not exceeding 5000 words) discuss current trends in research in a field that will be of interest to readers outside the field; interdisciplinary topics: science policy and administration; or some aspect of the application of science and technology to human needs or the impact of science and technology on society/ecosystems/life. They should include a summary not exceeding 100 words, introductory paragraph(s), brief subheads at appropriate places to point to what follows, illustrations that will help a general reader, and references.

Review articles (not exceeding 5000 words) are expected to survey and discuss recent developments in a field. They should be well focused and organized, and avoid a general, 'textbook' style.

Research articles (not exceeding 4000 words) should report research results of fairly major significance. They should include an abstract not exceeding 100 words, introductory paragraph(s), and brief subheads.

Research communications (not exceeding 2000 words) should contain important findings that are novel and of fairly broad interest. They should include a brief abstract and an introductory paragraph. Text should not be broken up under subheads.

Correspondence includes letters, not exceeding 500 words, that are of general interest to scientists. All letters cannot be published.

Scientific correspondence contains technical comments, including those on articles or communications published in *Current Science* within the previous six months. Letters may be reviewed and edited.

Research news articles are intended to inform nonspecialists about recently published advances or important findings discussed at a meeting. Authors should also send a copy of the paper(s) on which the article is based. Meeting reports should avoid merely listing brief accounts of topics discussed, and must convey to readers the significance of an important advance.

Research accounts articles are intended to be personalized reviews of research from the authors' own laboratory, based on a body of published work. The articles must provide appropriate background to the area in a concise introduction, which should also serve to place the author's work in proper perspective. Articles will normally

not exceed 8 to 10 printed pages.

Opinion articles present views on issues related to science and scientific activity. **Commentary articles** should contain expository notes on issues related to science and scientific activity.

Book reviews. Unsolicited reviews will also be considered. Reviews that merely 'list' brief descriptions of the contents cannot be published. Reviews should have 'context' and convey some information about the subject of the book.

Historical commentary and notes inform readers about interesting aspects of personalities or institutions of science or about watershed events in the history/development of science; most are expected to relate to India. Illustrations are welcome. Brief items will also be considered.

MANUSCRIPT PREPARATION

Manuscripts should be typed double-spaced on one side of white bond paper (21 × 28 cm). The pages should be numbered consecutively, starting with the title page and through the text, reference list, tables and figure legends. The title should be brief, specific and amenable to indexing. Not more than five **keywords** should be indicated separately; these should be chosen carefully and must not be phrases of several words. **Summary and abstract** should not have more than 100 words and should convey the main point of the paper, *outline the results and conclusions, and explain the significance of the results.*

Text. All papers should have a brief introduction. The text should be intelligible to readers in different disciplines and technical terms should be defined. Tables and figures should be referred to in numerical order. All **symbols and abbreviations** must be defined, and used only when absolutely necessary. Superscripts and subscripts and ambiguous characters should be clearly indicated. **Units of measure** should be metric or, preferably, SI. Methods should, as far as possible, be described briefly in appropriate table and figure legends.

Figures. In the case of line drawings one set of originals (without any lettering) is sufficient, with two copies containing lettering. In the case of photographs good prints are required with each copy of the manuscript; photocopies are not acceptable. Line drawings should be roughly twice the final printed size. The correct orientation should be indicated if not clear.

Photomicrographs and other photographs that require it must have a scale bar, which should be defined clearly in the legend. Primary data should be submitted as far as possible (e.g. actual photographs of electrophoretic gels rather than idealized diagrams).

References should be numbered in the order in which they appear, first through the text and then through table and figure legends. The following are examples of ways of writing references.

1. Mukundan, T. and Kishore, K., *Curr. Sci.*, 1991, 60, 355-362.
2. Constantine, G., in *Biology of Bats* (ed. Wimsatt, W. A), Academic Press, New York, 1970. vol 1, pp. 319-322.

Acknowledgements should be brief. Footnotes are not allowed except to identify the corresponding author if not the first.

Cover photographs. Good photographs (colour or black and white) that pertain to a submitted paper will be considered for use on the cover. Good *prints* and a legend should be submitted with the manuscript. In the case of a colour picture, a transparency will be required for printing if accepted.

PROOFS AND PUBLICATION

Two sets of galley proofs are sent to the corresponding author. A reprint order form accompanies the proofs.

object?) determines the relative differential hemispheric activation⁴.

Shaywitz and his colleagues⁵ asked their subjects (19 males and 19 females) to map the given letter strings onto phonological representations. In other words, the subjects are to determine if two nonsenseword strings rhymed. The test involves phoneme coding. Neural activation during rhyming in males was lateralized predominantly to the left inferior frontal regions (mean number of pixels activated were 11.7 and 5.0 for the left hemisphere and the right hemisphere respectively). In contrast, activation during this same task in females engaged this region bilaterally (mean number of corresponding number of pixels activated for females were 9.4 and 12 in the left hemisphere and the right hemisphere respectively). Shaywitz and his associates thus demonstrated that in a site uniquely serving phonological processing, inferior frontal gyrus (IFG), females devote greater right hemispheric resources to the task. Also, cumulatively larger areas in IFG in both the hemispheres are seen activated in females when compared to males.

However, Fink and his associates in their experiments apparently used only male subjects. Will the results from the female subjects be different in any manner for global/local processing of letter-based and object-based stimuli? In other words, who among the males and females are more likely to 'miss the wood for the trees'? Does Shaywitz *et al.*'s work indicate a natural flair for lullaby among females? I am sure the evolutionists among the biologists are sure to hijack the answers to these and similar questions (as and when available) to see in the answers some biological advantage to the *Homo sapiens* as a species! In the meantime, can we not think of enlarging the scope of our endeavours in brain research so as to find applications in as broad a realm as education, besides the efforts directed towards the amelioration of disorders/diseases afflicting the brain⁶?

1. Springer, S. P. and Deutsch, D., *Left Brain, Right Brain*, W. H. Freeman, New York, 1983, 4th ed.
2. Albert, S. M., *Science*, 1997, **275**, 1547.
3. Fink, G. R., Halligan, P. W., Marshall, J. C., Frith, C. D., Frackowiak, R. S. J. and Dolan, R. J., *Nature*, 1996, **382**, 626-628.

4. Fink, G. R., Marshall, J. C., Halligan, P. W., Frith, C. D., Frackowiak, R. S. J. and Dolan, R. J., *Proc. R. Soc. London*, 1997, **B264**, 487-494.
5. Shaywitz, B. A., Shaywitz, S. E., Pugh, K. R., Todd Constable, R., Skudlarski, P., Fulbright, R. K., Bronen, R. A., Fletcher, J. M., Shankweiler, D. P., Katz, L. and Gore, J. C., *Nature*, 1995, **373**, 607-609.
6. Ramakrishna, T., *Curr. Sci.*, 1997, **73**, 1045-1047.

ACKNOWLEDGEMENTS. I thank Prof. P. Balaram, Chairman, MBU, IISc., for kindly suggesting that I write this article. I am greatly beholden to Prof. K. R. K. Easwaran, MBU, IISc., and Prof. C. N. R. Rao, President, Jawaharlal Nehru Centre for Advanced Scientific Research, Bangalore for the kind invitation to be visiting scientist.

T. Ramakrishna was in the Department of Life Sciences, University of Calicut, Calicut 673 635, India. He is currently in the Jawaharlal Nehru Centre for Advanced Scientific Research and Molecular Biophysics Unit, Indian Institute of Science, Bangalore 560 012, India.

Magnetic reconnection on the Sun: Ultraviolet observations of bi-directional plasma jets

B. N. Dwivedi and Anita Mohan

Magnetic field reconnection transcends the traditional disciplines of laboratory, space and cosmic plasma physics. The process is known to occur in fusion devices such as tokamaks where it causes major disruption of the plasma confinement. It is thought to occur in solar flares and in other energetic events on the Sun. It takes place at planetary magnetopauses, and in planetary, as well as cometary, magnetic tails. Reconnection also plays an important role in other cosmic objects such as accretion discs, and in a variety of current sheets occurring in interplanetary, interstellar, and intergalactic space. Detailed study of reconnection processes in the laboratory, in computer simulations, in the Earth's magnetosphere and in the solar plasma is, therefore, in the forefront of current researches.

In cosmic plasmas, large scale lengths

L , large velocities V , and small electrical resistivities η combine to form large values of the magnetic Reynolds number $R_m = \mu_0 VL/\eta$, a condition in which the plasma and magnetic fields are tightly coupled or 'frozen' together. If the ratio of plasma to magnetic energy density is large, non-uniform motions in such plasmas often stretch magnetic loops or push differentially magnetized regions together into configurations, where magnetic field exhibits large shear, i.e. it changes direction and magnitude rapidly across a narrow electric current sheet. If the ratio is small, the magnetic field organizes the plasma motion instead. And the currents have a tendency to flow along magnetic field lines, a situation that also leads to sheared magnetic fields. All such field configurations contain free magnetic energy. Magnetic reconnection provides

a means of converting some or all of this energy to plasma kinetic and thermal energy and thus of changing the field configuration into a thermal equilibrium.

Magnetic reconnection is the process by which magnetic lines of force break and rejoin into a lower-energy configuration. This is considered to be the fundamental process by which magnetic energy is converted into plasma kinetic energy. The Sun has a large reservoir of magnetic energy. The energy released by magnetic reconnection has been invoked to explain both large-scale events, such as solar flares and coronal mass ejections, and small-scale phenomena, such as the coronal and chromospheric microflares that are likely candidates to heat and accelerate the solar wind.

The solar network (chromospheric and photospheric features arranged in a cel-