

Solubilization of inorganic phosphates by *Azospirillum halopraeferans*

Bacteria of the genus *Azospirillum* are widely distributed in the rhizosphere of tropical and subtropical grasses and sugarcane¹. *Azospirillum halopraeferans* is reported to occur in the rhizoplane of plants growing in saline soil in Brazil². Phosphates, widely distributed in nature in both organic and inorganic forms, are not readily available to plants in a bound state³. Many soil bacteria are reported to solubilize these insoluble phosphates through various processes⁴⁻⁷. A few reports have also indicated the P-solubilizing activity of some nitrogen fixers⁸⁻¹¹. This paper reports the P-solubilizing capacity of three different strains of *A. halopraeferans*.

Type strains (LMG7107, LMG 7108 and LMG 7109) obtained from J. Dobreiner, EMBRAPA, Brazil were used in this study. Individual strains were grown on the Sperber's medium¹² and Pikovskaya's medium¹³ in petri dishes for 3 days at $28 \pm 2^\circ\text{C}$. The size of the clear zones around the colonies showing solubilization of phosphate on incubation was noted. The results are expressed as solubilization efficiency ($E = \text{Solubilization diameter } (S) / \text{Growth diameter } (G) \times 100$) (ref. 14). For broth culture studies, single colonies grown on nutrient agar were inoculated into both the liquid media containing water insoluble P-sources. All the experiments were conducted in triplicate and the cultures were maintained for 16 days. Phosphate was determined by the paramolybdate blue method¹⁵. For analysis, the cultures were harvested on every alternate day, centrifuged at 10,000 rpm for 15 min and then subjected for estimation. pH of the medium was also recorded simultaneously.

While the strains showed weak zone of solubilization on Sperber's medium (Table 1, Figure 1), no activity was observed in Pikovskaya's medium in plate

assay. However, the liquid environment offered encouraging results where the three strains showed good activity in both the broths employed. Morris and Allen¹⁶, while studying the oxalate metabolism by micro-organisms echoed a conforming trend, where the organisms do not show clearing zone in plates and there was good calcium oxalate metabolism in liquid broth, attributed to simple cation dissociation. Periodical estimates of P in media revealed the potential of all the strains in releasing Pi from insoluble P sources. On Sperber's medium, the P concentration increased slowly, reached a peak on the tenth day and declined slowly on later days. Whereas in Pikovskaya's medium, the P concentration increased gradually achieving a peak

on the sixth day. Among the strains, LMG 7109 was good in Sperber's medium and solubilized $122.54 \mu\text{g P/ml}$, while LMG 7107 and LMG 7108 solubilized 84.61 and $47.19 \mu\text{g P/ml}$, respectively (Figure 2). In Pikovskaya's medium, LMG 7109 showed maximum solubilization ($1453.69 \mu\text{g P/ml}$) on 14th day, whereas in LMG 7107 and LMG 7108 it was 1379.6 and $1452.04 \mu\text{g P/ml}$ on 6th day itself (Figure 3). Statistically no significant difference was observed in P-solubilization among strains in Sperber's medium except between 8 and 12 days (Figure 2), but in Pikovskaya's medium the strains differed significantly on all days (Figure 3).

In general, the bacterial activity was slow initially, which gradually increased




Figure 1. Plate assay for P-solubilization by *A. halopraeferans*. *a*, Solubilization zone formed in Sperber's medium; *b*, Colonies without any halo zone in Pikovskaya's medium.




Figure 2. Change of P concentration in Sperber's medium.

Table 1. Solubilization efficiency (*E*) of *A. halopraeferans* strains in plate assay

Strain	Sperber's medium*	Pikovskaya's medium
LMG 7107	151.14	Nil
LMG 7108	151.69	Nil
LMG 7109	150.56	Nil

*Results presented are average of three replicates.

in the middle and declined later. Decrease in P concentration during initial stages in Sperber's medium (Figure 2) can be attributed to the utilization of existing P for growth and development of the organism⁶, in a later phase the bacteria would have started acting on the substrate for

want of nutrients, thus releasing P from insoluble sources. Or, the cells after the initial shock could have utilized the available free P for metabolism and later to acclimatize to the given environment or due to substrate stress they would have solubilizing P. The better performance of

bacteria in Pikovskaya's medium by all strains indicates the role played by the substrate to trigger the microbial action^{7,17}. The P concentration in solution did not follow a sigmoid curve type, but some fluctuations and then increase in P concentration during later days in LMG 7108 and LMG 7109, which may be due to cell lysis and P precipitation brought about by organic metabolites^{7,18,19}.

There are several potential mechanisms reported for phosphate solubilization, that include modification of pH by secretion of organic acids and protons or cation dissociation^{7,20-24}. In this study, pH values increased in Sperber's medium during early days and on incubation went down slowly (Figures 4 and 5). Maximum pH reduction recorded was 0.30 units in Sperber's medium and 0.53 units in Pikovskaya's medium. pH reduction was gradual, where it slowly went down and no revival was observed in later days. Results similar to this were observed by other workers^{25,26}, while studying cyanobacterium mediated P-solubilization.

A. halopraeferans, a non-glucose utilizing bacteria does not exhibit acidity in the presence of glucose⁴. Results obtained in the present study on glucose amended media indicate that acid production is not the only reason for P release into the media^{7,27,28} and this can be related to the cation dissociation process¹⁶. From the above results, it can be concluded that an interesting phenomenon of P-solubilization by nitrogen fixing *A. halopraeferans* could be further studied to find out the mechanism of action or the involvement of genes. A study on the molecular mechanism would throw light on the *ps* (phosphate solubilizing) genes that could be incorporated in agriculture as *A. halopraeferans* offers traits for nitrogen fixation, phosphate solubilization and salinity tolerance.




Figure 3. Change of P concentration in Pikovskaya's medium.




Figure 4. Change of pH in Sperber's medium.




Figure 5. Change of pH in Pikovskaya's medium.

1. Dobereiner, J. and Day, J. M., Proc. of 1st Int. Symp. on Nitrogen Fixation (eds. W. E. Newton and C. J. Nyman), Washington State University Press, 1976, p. 518.
2. Reinhold, B., Hurek, T., Baldani, I. and Dobereiner, J., in *Azospirillum IV Genetics, Physiology, Ecology* (ed. Klingmuller, W.), Springer-Verlag, New York, 1988, pp. 234-241.
3. Hayman, D. S., in *Soil Microbiology*, Butterworths, London, 1975, pp. 67-92.
4. Sperber, J. I., *Aust J. Agric. Res.*, 1958, 9, 778-781.

5. Banik, S. and Dey, P. K., *Plant Soil*, 1982, **69**, 353–364.
6. Molla, M. A. Z., Choudhary, A. A., Islam, A. and Huque, A., *Plant Soil*, 1984, **78**, 393–399.
7. Illmer, P. and Schinner, F., *Soil Biol. Biochem.*, 1992, **24**, 389–395.
8. Will, M. E. and Sylvia, M., *Appl. Environ. Microbiol.*, 1990, **56**, 2073–2079.
9. Halder, A. K., Mishra, A. K. and Chakrabarthy, P. K., *Indian J. Microbiol.*, 1990, **30**, 311–314.
10. Halder, A. K., Mishra, A. K. and Chakrabarthy, P. K., *Indian J. Exp. Biol.*, 1991, **29**, 28–31.
11. Maheshkumar, K. S., Krishnaraj, P. U. and Algawadi, A. R., *Curr. Sci.*, 1999, **76**, 874–875.
12. Sperber, J. I., *Nature*, 1957, **180**, 994.
13. Pikovskaya, R. E., *Mikrobiologiya*, 1948, **17**, 362.
14. Nguyen, C., Yan, W., Le Tacon, F. and Lapayrie, F., *Plant Soil*, 1992, **143**, 193–199.
15. Olsen, S. R. and Sommers, L. E., in *Methods in Soil Analysis* (eds Page, A. L., Miller, R. H. and Keeney, D. R.), 1982, pp. 403–430.
16. Morris, S. J. and Allen, M. F., *Biol. Fertil. Soils*, 1994, **18**, 255–259.
17. Nahas, E., *World J. Microbiol. Biotechnol.*, 1996, **12**, 567–572.
18. Khan, J. A. and Bhatnagar, R. M., *Fertil. Technol.*, 1977, **14**, 329–333.
19. Babenko, Yu. S., Tyrygina, G. I., Grigor'ev, E. F., Dohgikh, L. M. and Borisowa, T. I., *Mikrobiologiya*, 1984, **53**, 533–539.
20. Louw, H. A. and Webley, D. M., *J. Appl. Bacteriol.*, 1959, **22**, 227–233.
21. Katzelson, H., Peterson, E. A. and Rouatt, J. W., *Can. J. Bot.*, 1962, **40**, 1181.
22. Cunningham, J. E. and Kuiack, C., *Appl. Environ. Microbiol.*, 1992, **58**, 1451–1458.
23. Wenzel, C. L., Ashford, A. E. and Sumner, B. A., *New Phytol.*, 1994, **128**, 487–496.
24. Seshadri, S., Ph D thesis, Bharathidasan University, Trichy, India, 1995.
25. Bose, P., Nagpal, U. S., Venkataraman, G. S. and Goyal, S. K., *Curr. Sci.*, 1989, **58**, 165–166.
26. Roychoudhury, P. and Kaushik, B. D., *Curr. Sci.*, 1989, **58**, 569–570.
27. Asea, P. E. A., Kucey, R. M. N. and Stewart, J. W. B., *Soil Biol. Biochem.*, 1988, **20**, 459–464.
28. Salih, H. M., Yahya, A. I., Abdul-Rahem, A. M. and Munam, B. H., *Plant. Soil*, 1989, **120**, 181–185.

ACKNOWLEDGEMENT. R.M. thanks Dr Johanna Dobereiner, EMBRAPA, Brazil for providing type strains.

Received 23 March 2000; revised accepted 5 July 2000

S. SESHADRI^{*,‡}
R. MUTHUKUMARASAMY[#]
C. LAKSHMINARASIMHAN[†]
S. IGNACIMUTHU^{*}

^{*}Entomology Research Institute,
Loyola College,
Chennai 600 034, India

[#]Main Bio-control Research Laboratory,
Tamil Nadu Co-operative Sugar
Federation,
Chengalpattu 603 001, India

[‡]Department of Botany,
AVVM Sri Pushpam College,
Poondi 613 503, India